

Alphabetical Index Changes (for use in South Africa)

Changes made to the 2008 Edition up to and including January 2013 of the WHO Corrigenda.

The changes highlighted in blue may not be reflected in the 2010 edition of ICD-10 (also referred to as the 4th edition of ICD-10).

South African Implementation date: 01 June 2013

Volume 3

ALPHABETIC INDEX

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Revise code	Introduction
	General arrangement of the Index
	Multiple diagnoses
	The Tabular List includes a number of categories for the classification of two or more conditions jointly reported, e.g. "Influenza with
	pneumonia" (J11.0), "Acute appendicitis with generalized peritonitis" (K35.0K35.2). Such combinations of conditions, which are specifically
	classified in the Tabular List, also appear in the Index.
	Abnormal, abnormality, abnormalities — see also Anomaly
Add subterm:	- karyotype R89.8
Add subterm:	- kidney function test R94.4
	- labour NEC O75.8
Revise codes	Abnormal, abnormalities – see also Anomaly
	- course, eustachian tube Q17.8 Q16.4
	Abortion(complete) (incomplete) O06
	- attempted (failed) (induced) (nonmedical) O07.9
	complicated by
	sepsis O07.5 septic shock O07.5
Revise subterm:	septic snock 007.5 septicemia 007.5
Kevise subtefill.	Septicemia Oo7.3

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	medical complicated by
	sepsis O07.0 ——septicemia O07.0
	Abortion -see list of fourth characters to be used with categories O03-O06 - complicated (by)
	sepsis .5 .0 .0 septic shock .5 .0 .0 - septicemia .5 .0 .0
Add dagger symbols and asterisk codes	Abscess (embolic) (infective) (metastatic) (multiple) (pyogenic) (septic) L02.9 - amebic A06.4 brain (and liver or lung abscess) A06.6† G07* liver (without mention of brain or lung abscess) A06.4† K77.0*
Revise code	 - lung (and liver) (without mention of brain abscess) A06.5† J99.8* - specified site NEC A06.8 - spleen A06.8† D77*
	 bone (subperiosteal) M86.8 M86.9 spleen D73.3 amebic A06.8 † D77*
Revise codes at subterms	Abscess (embolic)(infective)(metastatic)(multiple)(pyogenic)(septic) L02.9 appendix K35.1K35.3
	- ileocecal <u>K35.1K35.3</u> periappendicular <u>K35.1K35.3</u>
	- peritoneum, peritoneal (perforated)(ruptured) K65.0 - with appendicitis K35.1 K35.3
Revise code	- vermiform appendix <u>K35.1</u> <u>K35.3</u> Absorption

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	 chemical T65.9 specified chemical or substance (see Table of drugs and chemicals) through placenta (fetus or newborn) P04.8 P04.9 suspected, affecting management of pregnancy O35.8
Revise lead term, subterm and code	Accreta placenta (with hemorrhage) O72.0 — without hemorrhage O73.0 — complication during delivery (with hemorrhage) O72.0 — without hemorrhage O73.0 — maternal care O43.2
Add subterms	Acquired–see also condition – immunodeficiency syndrome (AIDS)(see also Human immunodeficiency virus (HIV) disease) B24 – complicating pregnancy, childbirth or the puerperium O98.7
Revise code	Adenofibroma (M9013/0) — see also Neoplasm, benign - prostate N40 D29.1
Revise code	Adenoma (M8140/0) — see also Neoplasm, benign - prostate N40-D29.1 Adenocarcinoma (M8140/3) — see also Neoplasm, malignant
Add subterm:	- mucinous (M8480/3) metastatic (M8480/6) - see Neoplasm, secondary
Revise code	Adenomyoma (atypical polypoid) (M8932/0) — see also Neoplasm, benign - prostate N40-D29.1
Revise subterms	Adherent –see also Adhesions – placenta (with hemorrhage)(morbidly) O72.0 O43.2 — without hemorrhage O73.0
Revise and add subterms	Adhesions, adhesive (postinfective) K66.0 - meninges (cerebral) (spinal) G96.1 - cerebral G96.1 - congenital Q04.8 - congenital Q07.8 - spinal G96.1 - congenital Q06.8 - tuberculous (cerebral)(spinal)A17.0† G01*
Add subterm	AIDS (related complex)(see also Human, immunodeficiency virus (HIV) disease) B24 – complicating pregnancy, childbirth or the puerperium O98.7
Delete subterm	Albuminuria, albuminuric (acute) (chronic) (subacute) (see also Proteinuria) R80 - complicating pregnancy, childbirth or puerperium O12.1

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	with
	— gestational hypertension (see also Pre eclampsia) 014.9 — pre existing hypertensive disorder 011
	- gestational O12.1
	with
	— gestational hypertension (see also Pre eclampsia) O14.9
	— pre existing hypertensive disorder O11
	- orthostatic N39.2
	- postural N39.2
	- pre eclamptic (see also Pre eclampsia) O14.9
	affecting fetus or newborn P00.0 severe O14.1
	— affecting fetus or newborn P00.0
	Alcohol, alcohol- induced
	- intoxication (acute) F10.0
	with
	delirium F10.0
Delete subterm	— dependence F10.2
	Alcohol, alcoholic, alcohol induced
	detoxification therapy Z50.2dependence F10.2
Revise subterms	- intoxication (acute) F10.0
	- with delirium F10.0
	— delirium F10.0
Add new lead	Amianthosis J61
term	
	Anaphylactoid shock or reaction – see Shock, anaphylactic
Add lead term and	Anaphylactoid syndrome of pregnancy O88.1
add code:	Anaphylaxis T78.2
Revise code	Anastomosis
	- intestinal K63.8 <u>Z98.0</u>
	 - complicated NEC K91.8 involving urinary tract N99.8
	Anemia D64.9
	Autina Dut.7
Revise and add	- in neoplastic disease NEC (M8000/1) (see also Neoplasm) D48.9† D63.0*

ALPHABETIC INDEX ENTRIES
chronic kidney disease
stage 3 N18.3† D63.8* stage 4 N18.4† D63.8*
stage 5 N18.5† D63.8
unspecified N18.9† D63.8*
neoplastic disease NEC (M8000/1) (see also Neoplasm) D48.9† D63.0*
- infantile D64.9
Anemia D64.9
- myelodysplastic (M9989/1)(related to alkylating agent) (related to Epipodophyllotoxin) (related to therapy) D46.9
– myelofibrosis (M9960/1) D47.1† D63.0*
 - myeloproliferative (M9960/1) D47.1
- refractory (M9980/1)(related to alkylating agent) (related to Epipodophyllotoxin) (related to therapy) NEC D46.4
with
<u>– – dysplasia, multi lineage D46.5</u>
excess of blasts (M9983/1) D46.2
with blasts in transformation (M9984/1) D46.3 C92.0
<u></u> of blasts (RAEB I)(RAEB II) D46.2 hemochromatosis (M9982/1) D46.1
sideroblasts, ring ed (M9982/1) D46.1
- sideroblastic (M9982/1) D46.1
– without sideroblasts, ringed (M9981/1) D46.0
Anesthesia, anesthetic (see also Effect, adverse, anesthesia) R20.0
- death from
correct substance properly administered T88.2
during delivery O74.8 O74.9
in pregnancy O29.9
Aneurysm (anastomotic) (artery) (cirsoid) (diffuse) (false) (fusiform) (multiple) (saccular) I72.9
- arteriovenous (congenital)(peripheral) Q27.3
acquired I77.0
brain I67.1 pulmonary I28.0
brain Q28.2
ruptured I60.8

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Revise and add and subterms:	precerebral vessels (nonruptured) Q28.0 ruptured I72.5 specified site NEC Q27.3 acquired I77.0 traumatic (complication)(early) T14.5 dissecting (see also Dissection, artery) I72.9 aorta (any part)(ruptured) I71.0 syphilitic A52.0† I79.0* precerebral-congenital (nonruptured) Q28.1 NEC I72.5 acquired (ruptured) I72.5 carotid (internal) I72.0 vertebral I72.5 congenital (nonruptured) Q28.1 - pulmonary I28.1
Revise code and add subterms	Aneurysm - brain I67.1 - congenital Q28.3 berry (nonruptured) Q28.3 ruptured I60.7 ruptured I60.7
Revise and add subterms	Aneurysm (anastomotic)(artery)(cirsoid)(diffuse)(false)(fusiform)(multiple) (saccular) I72.9 - arteriovenous (congenital)(peripheral) Q27.3 - acquired I77.0 - brain Q28.2 ruptured I60.8 - precerebral vessels (nonruptured) Q28.0 ruptured NEC I72.5 - precerebral NEC I72.5 - acquired (ruptured) I72.5 - acquired (ruptured) I72.5

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	carotid (internal) I72.0
5.1	vertebral I72.5I72.6
Delete morphology code and revise code	Angioendotheliomatosis (M9712/3) C85.79
Revise codes	Anomaly, anomalous (congenital)(unspecified type) Q89.9
	– eustachian tube Q17.8 <u>Q16.4</u>
	Anthrax A22.9
	- respiratory A22.1
Revise subterm:	- - septicaemia sepsis A22.7 - specified manifestation NEC A22.8
A 1.1 1	Anthrax A22.9
Add dagger symbol and	- cerebral A22.8 <u>† G01*</u>
asterisk codes	- gastrointestinal A22.2 † K93.8*
asterisk codes	- inhalation A22.1† J17.0*
	– pulmonary A22.1 † J17.0*
	– respiratory A22.1 <u>†J 17.0*</u>
Add lead term and code	Anticardiolipin syndrome D68.6
Add lead term and	Antiphospholipid syndrome D68.6
code	
Revise code	Antithrombinemia (see also Circulating anticoagulants) D68.3 D68.5
	Apnea, apneic (spells) R06.8
Add non-essential	- newborn NEC (obstructive) P28.4
modifiers and	sleep (central) (primary) P28.3
subterm	- prematurity (of) P28.4 - sleep (central) (obstructive) G47.3
Revise code:	Apoplexia, apoplectic I64
Revise code.	
	- uremic N18. 85 † I68.8*
	Appendicitis K37
	- with
Revise subterms:	— perforation or rupture K35.0
	peritoneal abscess K35.1 <u>K35.3</u>
	<u> peritonitis, generalized K35.2</u>

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	with mention of perforation or rupture K35.2 peritonitis, localized K35.9 K35.3 with mention of perforation or rupture K35.0 K35.3 generalized K35.0 acute (catarrhal) (fulminating) (gangrenous) (obstructive) (retrocecal) (suppurative) K35.9 K35.8 with perforation or rupture K35.0 peritoneal abscess K35.1 K35.3 peritonitis, generalized K35.2 with mention of perforation or rupture K35.2 peritonitis, localized K35.9 K35.3
	with mention of perforation or rupture K35.0 K35.3 generalized-K35.0 - amebic A06.8
Revise subterms	Appendicitis K37 - with peritonitis, generalized NECK35.2K35.3 generalized (with perforation or rupture) K35.2 with mention of perforation or rupture K35.2 localized (with perforation or rupture K35.3 peritonitis, localized K35.3 with mention of perforation or rupture K35.3 with mention of perforation or rupture K35.3 acute (catarrhal) (fulminating) (gangrenous) (obstructive) (retrocecal) (suppurative) K35.8 with peritonitis, generalized NECK35.2K35.3 with mention of perforation or rupture K35.2 generalized (with perforation or rupture) K35.3 peritonitis, localized (with perforation or rupture) K35.3 peritonitis, localized K35.3 with mention of perforation or rupture K35.3
Add subterm	ARC (AIDS-related complex) (see also Human, immunodeficiency virus (HIV) disease) B24 - complicating pregnancy, childbirth or the puerperium O98.7
Add modifier at subterm	Arteriosclerosis, arteriosclerotic (diffuse) (disease) (general) (obliterans) (senile) (with calcification) I70.9 - mesenteric (chronic) K55.1

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Add index subterm	Arteriosclerosis, arteriosclerotic (diffuse) (disease) (general) (obliterans) (senile) (with calcification) I70.9 <u>- carotid artery I65.2</u>
Add subterms and revise codes	Arthritis, arthritic (acute) (chronic) (subacute) M13.9 - in (due to) - enteritis NEC A09.9† M01.8*M03*
Delete	infectious NEC A09.0† M01.8*M03* Arthritis, arthritic (acute)(chronic)(subacute) M13.9
morphology code	 in (due to) leukemia NEC (M9800/3) C95.9† M36.1* multiple myeloma (M9732/3) C90.0† M36.1* myelomatosis (multiple) (M9732/3) C90.0† M36.1* reticulosis, malignant (M9713/3) C85.7 C86.0† M14.8*
Add subterms and revised codes	Arthropathy (see also Arthritis) M13.9 in (due to) endocrine disorder NEC E34.9 M14.5* enteritis NEC A09.9† M03* infectious NEC A09.0† M03* enteropathic NEC K52.9+ A09.9+ M07.6*
	infectious A09.0+ M07.6*M03* Arthropathy (see also Arthritis) M13.9 - in (due to)
Add dagger/asterisk combinations	erythema multiforme L51† M14.8* nodosum L52† M14.8 hematologic disorders NEC (see also categories D50-D77) D75.9† M36.3*
Delete morphology codes	 - hepatitis viral (see also categories B15– B19) B19.9† M03.2* - leukemia NEC (M9800/3) (see also categories C91–C95) C95.9† M36.1* - malignant histiocytosis (M9720/3) C96.18† M36.1* - mycosis NEC (see also categories B35–B49) B49† M01.6* - viral hepatitis (postinfectious) NEC (see also categories B15–B19) B19.9† M03.2*
Revise code	Arthropathy (see also Arthritis) M13.9 - in (due to)

INSTRUCTION	ALPHABETIC INDEX ENTRIES
ranges	hematologic disorders NEC (see also categories D50 D77 D50-D62, D64-D76) D75.9† M36.3*
Delete	Arthropathy
morphology code	in (due to)
D : 1	multiple myeloma (M9732/3) C90.0† M36.1*
Revise code	Astereognosis R41.3-R48.1
Revise codes	Atresia, atretic
	overteekien tuho 017 9 016 4
Add subterms and	- eustachian tube Q17.8 Q16.4 Atrophy, atrophic - continued
codes	Act opiny, act opine – continueu
codes	- macular (dermatological) L90.8
	- multiple system (brain)(CNS) G90.3
	- muscle, muscular M62.5
	Atrophy, atrophic
Delete subterm	- kidney (senile) (terminal) (see also Sclerosis, renal) N26
Delete subterm	hydronephrotic N13.3
	with infection N13.6
Revise code	— with hypertension (see also Hypertension, kidney)I12 Attack
Revise code	- vasomotor R57.9- R55
Add lead terms	AVNRT (atrioventricular nodal re-entrant tachycardia) I47.1
Add icad terms	AVRT (atrioventricular nodal re-entrant tachycardia) I47.1
Revise subterm:	Bacteremia A49.9
	-with sepsis — see Septicaemia Sepsis
	Barrett's
	- disease K22.7
	- esophagus K22.7
Add subterm:	malignant - see Neoplasm, esophagus, malignant
	- syndrome K22.7
	- ulcer K22.1
Add lead term and	Bedfast, NEC R26.3
code, revise	- requiring health care provider Z74.0
subterm	

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Add parentheses	Bednar('s)
and revise cross reference	– aphthae K12.0
	– tumor (M8833/3) — see Neoplasm, <u>skin</u> , malignant
Revise	Bedsore L89
Add subterms	- stage
	<u>IL89.0</u>
	<u> II L89.1</u>
	<u> III L89.2</u>
	<u> IV L89.3</u>
Add lead term	Bleb(s) R23.8
and subterms	– emphysematous (lung) (solitary) J43.9
	<u>– endophthalmitis H59.8</u>
	<u>– inflammed (infected), postprocedural H59.8</u>
Add lead term	Blebitis, postprocedural H59.8
and subterms	
Revise subterm	Bleeding (see also Hemorrhage) R58
	- hemorrhoids NEC 184.8(– see Hemorrhoids)
Revise codes	Blindness (acquired)(congenital)(binocular)(both eyes)(congenital) H54.0
add subterms	
	- hysterical F44.6
	- mind R48.8
	- monocular H54.4 - night H53.6
	- vitamin A deficiency E50.5
	- one eye (other eye normal) H54.4
	- low vision, other eye H54.1 H54.4
Modify subterm	Block
, , , , , , , , , , , , , , , , , , ,	– kidney (see also Failure, renal-kidney N19)
	Blood
Add non-essential	- transfusion (session) Z51.3
modifier and	reaction or complication -see Complications, transfusion
code:	without reported diagnosis Z51.3

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	Brill–Symmers' disease (M9690/3) C82.9
Add subterms	Bronchiolitis (acute) (infective) (subacute) J21.9 – due to
	 – external agent (see Bronchitis, acute or subacute, due to) – human metapneumovirus J21.1 – respiratory syncytial virus J21.0
Add subterms	Bronchitis (diffuse) (fibrinous) (hypostatic) (infective) (membranous) (with tracheitis)(15 years of age and above) J40 – acute or subacute (with bronchospasm or obstruction) J20.9 – due to
	Haemophilus influenzae J20.1 human metapneumovirus J20.8
Revise code	Bronchitis - capillary (see also Pneumonia, broncho) <u>J18.0 J21.9</u>
	Brucella, brucellosis (infection) A23.9
Revise subterms:	- mixed A23.8 - septicaemia sepsis A23.9
Revise subterilis.	- septitaerinia sepsis A23.9 - melitensis A23.0 - specified NEC A23.8
Add lead term and	Burkholderia NEC A49.8
subterms	- mallei A24.0
	as the cause of disease classified elsewhere B96.8
	- pseudomallei (see also Melioidosis) A24.4 - as the cause of disease classified elsewhere B96.8
Revise lead term.	Burkitt's
and codes in	— cell leukemia (M9826/3) C91. 0 8
subterms. Delete	– lymphoma (malignant) (M9687/3) C83.7
morphology code	small noncleaved, diffuse (M9687/3) C83.7
	undifferentiated (M9687/3) C83.7
	- tumor (M9687/3) C83.7
	- type
	 – acute lymphoblastic leukemia (M9826/3) C91.08 – undifferentiated (M9687/3) C83.7
	Cachexia R64
Revise codes:	- cancerous (M8000/3) C80 C80
	– malignant (M8000/3) C80 <u>C80</u>

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	Calcification
Add subterm	- heart (see also Degeneration, myocardial) I51.5
	- reart (see diso Degeneration, myocardia) 151.5
	- idiopathic infantile arterial Q28.8
	- intervertebral cartilage or disk (postinfective) M51.8
Add subterm	Calculus
	honotohiliam: V90 9
Add subterms	- hepatobiliary K80.8 Cancer (M8000/3) — (see also Neoplasm, malignant)
Add subterills	Note: The term
	- non-responsive to antineoplastic drugs U85
	<u>- refractory U85</u>
	Candidiasis, candidal B37.9
Revise subterm:	- resulting from HIV disease B20
Revise subteriii:	- septicemia sepsis B37.7 Carcinoid (tumor) (M8240/3) – see also Neoplasm, malignant
Revise code:	– goblet cell (M8243/3) C80 <u>C80</u>
	Carcinomatosis
D . 1 1	'C 1 '- (MOO10/C) C00C70 0
Revise code and add subterms:	 unspecified site (M8010/6) C80C79.9 primary site not indicated C80.9
add subterilis.	- primary site unknown, so stated C80.0
	Carcinomatosis
Delete subterms	- unspecified site (M8010/6) C79.9
	— primary site not indicated C80.9 — primary site unknown, so stated C80.0
	Caries
	- bone NEC A18.0† M90.0*
	- cementum K02.2
Add subterms	arrested K02.3
Add subtellis	specified NEC K02.8
	- dental K02.9 with pulp exposure K02.5
	with pulp exposure K02.5

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	- tooth, teeth K02.9
	with pulp exposure K02.5 Cat
Add subterm	eye syndrome Q92.8
	Cesarean
	- emergency O82.1
	- operation or section NEC (see also Delivery, cesarean) O82.9 - with hysterectomy O82.2
	affecting fetus or newborn P03.4
	post mortem, affecting fetus or newborn P01.6
	preterm NEC O60.1
Add subterms	<u> with spontaneous labor O60.1</u>
	without spontaneous labor O60.3
	previous, affecting management of pregnancy O34.2
Add dagger	Chagas'(Mazza) disease (chronic) B57.2 – with
symbol and asterisk code	- with - nervous system involvement B57.4† G99.8*
Add lead term and	Chairfast-NEC R26.3
code; revise	-requiring health care provider Z74.0
subterm	
	Charley-horse (quadriceps) S76.1
	- muscle, except quadriceps (see Sprain)
Revise subterms:	non traumatic - see Cramp(s)
	traumatic - see Injury, muscle
Add subterm	Checking (of) - cardiac pacemaker (battery) (electrode(s)) (pulse generator) Z45.0
	- device
	cardiac NEC Z45.0
	contraceptive Z30.5
	Chill(s) R68.8
	- with fever R50.8
Revise subterm:	-septic — see Septicaemia Sepsis
Delete	Chloroma (M9930/3) C92.3
morphology code	Ch. L. d. P. NIEC WOO. 1
Add index	Cholestasis NEC K83.1

INSTRUCTION	ALPHABETIC INDEX ENTRIES
subterms:	- complicating pregnancy, childbirth or the puerperium (intrahepatic) O26.6
	Chondrodysplasia Q78.9
Add subterm:	- metaphyseal (Jansen's) (McKusick's) (Schmid's) Q78.5
Add dagger	Coccidioidomycosis, coccidioidosis B38.9
symbol and	– pulmonary B38.2 <u>†J99.8* J17.2*</u>
asterisk code	acute B38.0 <u>†</u> J99.8* <u>J17.2*</u>
	chronic B38.1 <u>†J99.8* J17.2*</u>
Revise modifiers	Colitis (acute) (catarrhal) (hemorrhagic) (presumed infectious) (see also Enteritis) A09.9
and add revised	- allergic K52.2
codes	- amebic (acute) (see also Amebiasis) A06.0
	nondysenteric A06.2
	- anthrax A22.2
	- bacillary (see also Infection, <i>Shigella</i>) A03.9
	- balantidial A07.0 - chronic (noninfective noninfectious) K52.9
	- coccidial A07.3
	- cystica superficialis K52.8
	- dietary counseling and surveillance (for) Z71.3
	- dietetic K52.2
	- due to radiation K52.0
	- food hypersensitivity K52.2
	- giardial A07.1
	- granulomatous K50.1
Add subterms and	- hemi-, left K51.5
code	- indeterminate K52.3
Revise code	- infectious (see also Enteritis, infectious) A09.0
	- ischemic K55.9
	acute (fulminant) (subacute) K55.0
	chronic K55.1
	fulminant (acute) K55.0
Add subterms	<u>- left sided K51.5</u>
	- noninfective noninfectious K52.9
Modify term	specified NEC K52.8
	- pan-, ulcerative (chronic) K51.0
	- polyposa K51.4
Delete text	-presumed noninfective K52.9
	- protozoal A07.9

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	- pseudomembranous A04.7 - regional K50.1
Revise code	- septic (<i>see also</i> Enteritis, infectious) <u>A09.0</u> - spastic K58.9
	- spastic K38.9 - with diarrhea K58.0
	- toxic K52.1
	- trichomonal A07.8
	- tuberculous (ulcerative) A18.3+ K93.0*
	- ulcerative (chronic) K51.9 - specified NEC K51.8
	Colitis (acute)(catarrhal)(hemorrhagic) (see also Enteritis A09.9)
	- coccidial A07.3
Add subterms	– <u>collagenous K52.8</u>
	- left sided K51.5
	- lymphocytic K52.8
	- microscopic K52.8
	- noninfectious K52.9
A 11 1.	specified NEC K52.8 Colitis (acute) (catarrhal) (hemorrhagic) (see also Enteritis) A09.9
Add subterm	(catarria) (nemorriagic) (see also Eliterius) A09.9
	- dietetic K52.2
	- drug-induced K52.1
	- due to radiation K52.0
Add subterms and	Colitis (acute) (catarrhal) (hemorrhagic) (see also Enteritis) A09.9
modify subterm	- granulomatous K50.1
	- infectious (see also Enteritis, infectious) A09.0
	due to Clostridium difficile A04.7
	- toxic NEC K52.1
	due to Clostridium difficile A04.7
Revise	Colitis(acute) (catarrhal) (hemorrhagic) (see also Enteritis, and note at category A09) A09.9
instructional note	

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Revise code	Coma R40.29
	– eclamptic (see also Eclampsia) R56.8O15.9
	Complications (from)(of)
	- accidental puncture or laceration during procedure T81.2
	- procedure (surgical or medical care) T88.9
Delete note:	Note: Whenever a complication of a procedure is not indexed or is not a synonym of an inclusion or indexed term, proceed as follows:
	Code to T80 T88:
	• early complications;
	mechanical complications
	Code to appropriate system chapter:
	• late complications
	• functional complications;
	- prosthetic device, graft or implant T85.9 Complications (from) (of)
	- orthopedic M79.9
Revise codes and	external device or appliance T88.8
subterms	internal device, implant or graft T84.9
	<u>— — infection or inflammation T84.5–T84.7</u> — — mechanical T84.3 T84.0–T84.4
	specified NEC T84.8
Add lead term	Complications (from)(of)
and subterms	- eye H57.9
	- – postoperative H59.9 - – blebitis H59.8
	Compression
Add lead term	-vena cava (inferior)(superior) I87.1 Compromised immune (system) NEC D89.9
Add Icad Icilii	Compulsion, compulsive
	Compression
Revise code and	- laryngeal nerve, recurrent <u>J38.7G52.2</u>

INSTRUCTION	ALPHABETIC INDEX ENTRIES
add subterm:	with paralysis of vocal cords and larynx J38.0
	Conditions arising in the perinatal period
	- depression
Revise code:	respiration, respiratory, newborn <u>P28.8P28.5</u>
Add subterms	Condition, conditions (arising in the perinatal period) (see also Disease) – gastro–esophageal reflux in newborn P78.8
	- gastro-esophageal reflux in newborn P78.8 - neonatal esophageal reflux P78.8
Modify index	Condition, conditions (arising in the perinatal period) (see also Disease)
subterms	- hepatitis
Sasterins	fetus or newborn (giant cell) (idiopathic) P59.2
	in toxoplasmosis, congenital (active) P37.1† K77.0*
	neonatal (giant cell) (idiopathic) (toxic) P59.2
	Constriction — see also Stricture
D	- larynx J38.6
Revise code:	congenital Q31.9 <u>8</u>
	Contraction, contracture, contracted - finger NEC M20.0
Revise code:	congenital Q68. 8 1
Revise code and	Cot death R95.9
add subterm	- with mention of autopsy R95.0
Revise code and	Crib death R95.9
add subterm	- with mention of autopsy R95.0
Revise	Cyst (colloid) (mucous) (retention) (simple)
	- congenital NEC Q89.8
	epiglottis Q31.8
	esophagus Q39.8
A 11 1.	fallopian tube Q50.4
Add subterms	<u> kidney Q61.0</u>
	larynx Q31.8
	liver Q44.7
Revise code and	- kidney (congenital <u>acquired</u>) Q61.0 <u>N28.1</u>
modifiers	- Ridney (congenital acquired) (consental A28.1 A28.1 Q61.0
	acquired congonian 1,2011 Q0110
Revise code and	- renal (congenital <u>acquired</u>) Q61.0 <u>N28.1</u>

INSTRUCTION	ALPHABETIC INDEX ENTRIES
modifiers	acquired congenital N28.1 Q61.0
	Cyst (colloid)(mucous)(retention)(simple)
Revise subterm:	- liver (idiopathic) K76.8
	congenital Q44.7
	Cyst (colloid)(mucous)(retention)(simple)
Add non-essential	– developmental K09.1
modifiers	odontogenic (glandular) K09.0
	- jaw (bone)(aneurysmal)(hemorrhagic)(traumatic) K09.2
	– – developmental (odontogenic)(<u>glandular</u>) K09.0
	ŭ i v
Add and revise	Cyst (colloid)(mucous)(retention)(simple) - congenital
subterms	- kidney Q61.0
	multicystic Q61.4
	polycystic Q61.3
	- kidney (acquired) N28.1
	congenital Q61.0
	calyceal (see also Hydronephrosis) N13.3
	multicystic (developmental) Q61.4
	<u>acquired N28.1</u>
	<u>polycystic multiple</u> Q61.3
	autosomal dominant (adult type) Q61.2
	autosomal recessive (infantile type) Q61.1
	- renal (acquired) N28.1
	congenital Q61.0
	multicystic Q61.4
	polycystic Q61.3
Revise subterms	Cystopyelitis (see also Pyelonephritis) N12
	<u>with calculus (impacted)(recurrent) N20.9</u>
	Death
Modify subterm	- instantaneous <u>unexplained</u> R96.0
	– sudden (cause unknown) <u>unexplained</u> R96.0

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	Death – obstetric (cause unknown) O95
	- affecting fetus or newborn P01.6
Revise codes:	 between 42 days and one year after delivery O96
	– one year or more after delivery O97 <u></u>
	Death
	- cot R95 <u>.9</u> with mention of autopsy R95.0
	with mention of autopsy R93.0 - crib R95.9
Revise code and	with mention of autopsy R95.0
add subterm	
	- sudden (cause unknown) R96.0
	infant R95 <u>.9</u>
	with mention of autopsy R95.0 Death
	- anesthetic
Revise codes	during delivery 074.8 074.9
110 / 150 0 0 000	in pregnancy O29.8 <u>O29.9</u>
	postpartum, puerperal O89.8
Revise	Decubitus (ulcer) L89
Add subterms	- cervix N86
	<u>- stage</u>
	<u>IL89.0</u>
	<u> II L89.1</u>
	<u> III L89.2</u>
- · · ·	IV L89.3
Revise code	Defect, defective - vision NEC <u>H54.7</u> <u>H54.9</u>
Add subterms	- VISION NEC <u>H34.7</u> H34.9 Deficiency
7 Rdd Subterinis	- antithrombin D68.5
	- protein (see also Malnutrition) E46
	<u> C D68.5</u>
	<u> S D68.5</u>

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	Deformity Q89.9
	- finger (acquired) M20.0
Revise codes:	congenital NEC Q68. <u>81</u>
	- thumb (acquired) M20.0
	congenital NEC Q68. <u>81</u>
	Deformity Q89.9
	flexion (joint) (acquired) M21.2congenital NEC Q68.8
	congenital NEC Qos.s hip or thigh (acquired) M21.2
Revise code	= - inp of thigh (acquired) W21.2 = congenital Q65.98
Revise codes	Deformity Q89.9
Tto vise codes	
	– eustachian tube (congenital) Q17.8 Q16.4
	Delay, delayed
	- development R62.9
Add subterm:	<u> global F89</u>
	intellectual (specific) F81.9
	Delirium, delirious (acute or subacute) (not alcohol- or drug-induced) F05.9
Add code:	- traumatic (see also Injury, intracranial) <u>S06.9</u>
	Delirium, delirious (acute or subacute) (not alcohol or drug induced) F05.9
Add subterm	– not superimposed on dementia F05.0
	<u>postoperative F05.8</u>
	– puerperal F05.8 Delivery (single) O80.9
	- complicated (by) O75.9
Revise and add	- complicated (by) 073.9 cervical dystocia (hypotonic) (failure of cervical dilatation) 062.0
subterms:	due to abnormality of cervix O65.5
Revise subterms	Delivery (single) O80.9
and codes	- complicated by
una codes	- adherent placenta O72.0 (morbidly) O43.2
	without hemorrhage O73.0
	placenta, placental
	accreta 072.0 043.2
	— without hemorrhage O73.0
	adherent (with hemorrhage) O72.0 (morbidly) O43.2

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	— without hemorrhage O73.0
	increta (with hemorrhage) O72.0 O43.2
	— without hemorrhage O73.0
	percreta O72.0 <u>O43.2</u>
	———— without hemorrhage O73.0
	Delivery (single) O80.9
	- complicated (by) O75.9
	compression of cord (umbilical) NEC O69.2
Add subterms	<u> around neck O69.1</u>
	contraction, contracted ring O62.4
	cord (umbilical)
	around neck, tightly or with compression O69.1
	<u> without compression O69.8</u>
	bruising O69.5
	complication O69.9
	specified NEC O69.8
	compression NEC O69.2
	entanglement O69.2
Add subterms	Delivery
	- cesarean
	<u> preterm NEC O60.1</u>
	<u> with spontaneous labor O60.1</u>
	<u> without spontaneous labor O60.3</u>
	fetus, fetal
	— prematurity O60.1
	- early onset (spontaneous) NEC O60 <u>.1</u>
	<u> with spontaneous labor O60.1</u>
	<u>-</u> - without spontaneous labor (cesarean section) (induction) O60.3
	- premature or preterm NEC O60.1
	<u> with spontaneous labour O60.1</u>
	without spontaneous labor (cesarean section) (induction) O60.3
	Delivery (single) O80.9
	- complicated (by) O75.9
Add and revise	laceration O70.9
subterms	<u>periurethral tissue O70.0</u>
	urethra O71.5
	<u> periurethral tissue O70.0</u>

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	vagina, vaginal wall (low)(minor)(deep) (high) (without perineal laceration)071.4-070.0 and muscles (perineal)(vaginal) 070.1
	and muscles (permear)(vagmar) 070.1 high (deep) (instrumental) (sulcus) 071.4
	— with perineum O70.0
	and muscles (perineal) (vaginal) O70.1
	Delivery (single) O80.9
	- complicated (by) O75.9 - laceration O70.9
Delete subterm	aceration 070.9 peritoneum 071.5
	periurethral tissue O70.0
	urethra O71.5
	— periurethral tissue O70.0
	Dementia (persisting) F03
D	
Revise code:	- uremic N18.85† F02.8*
Add subterms	Dementia (persisting) F03
	- degenerative (primary) F03
	- frontal lobe G31.0† F02.0*
	- frontotemporal G31.0† F02.0*
Add subterm	Dependence
	-on
	artificial heart Z99.4
	aspirator Z99.0
	Depression F32.9 -recurrent (<i>see also</i> Disorder, depressive, recurrent) F33.9
Add subterm:	- respiration, respiratory, newborn P28.5
rad subterm.	- respiratory centre G93.8
	Depression F32.9
Revise code	
	– bone marrow D75.8 <u>D61.9</u>
	Dermatitis L30.9
	- fungus G36.9
	specified type NEC B36.8

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Revise subterms	– gangrenosa, gangrenous <u>L88-infantum</u> <u>L08.0</u>
and codes:	— infantum R02
	Dermatofibrosarcoma (M8832/3) – <i>see also</i> Neoplasm, skin, malignant
Revise code:	– pigmented (M8833/3) C80 C80
Revised subterms	Dermatofibrosarcoma (protuberans) (M8832/3) - see also Neoplasm skin, malignant
and add non-	protuberans (M8832/3) C44.9
essential modifier	pigmented (M8833/3)-C80. see Neoplasm, skin, malignant
Delete	Di Guglielmo's disease (M9841/3) C94.0
morphology code	Diabetes, diabetic (mellitus)(controlled)(familial)(severe) E14.–
	Diabetes, diabetic (mellitus) (controlled) (familial) (severe) E14
Add subterm	
	- hemochromatosis E83.1
7	- hepatogenous E13
Revise modifiers	Diarrhea, diarrheal (disease) (endemie) (infantile) (presumed infectious) (summer) (see also Note at category A09) A09 A09.9
and add revised	- achlorhydric K31.8 - acute A09.9
codes	- acute A09.9 bloody A09.0
	- hemorrhagic A09.0
	- watery A09.0
	- allergic K52.2
	- amebic (see also Amebiasis) A06.0-
	with abscess - see Abscess, amebic
	acute A06.0
	chronic A06.1
	nondysenteric A06.2
	- and vomiting, epidemic A08.1
	- bacillary - see Dysentery, bacillary
	- balantidial A07.0
	- cachectic NEC K52.8
	- Chilomastix A07.8
	- chronic (noninfectivenoninfectious) K52.9 - coccidial A07.3
	- Coccidia A07.5
	- Cochin-China B78.0 - Dientamoeba A07.8
	- dietetic K52.2
	- due to

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	bacteria A04.9
	specified NEC A04.8 Campylobacter A04.5
	Clostridium perfringens (C) (F) A04.8
	Cryptosporidium A07.2
	Escherichia coli A04.4
	enteroaggregative A04.4
	enterohemorrhagic A04.3
	enteroinvasive A04.2
	enteropathogenic A04.0
	enterotoxigenic A04.1
	specified NEC A04.4
	food hypersensitivity K52.2
	specified organism NEC A08.5
	bacterial A04.8
	viral A08.3
	Staphylococcus A04.8
	virus (see also Eteritis, viral) A08.4
	Yersinia enterocolitica A04.6
	- dysenteric A09.0
Revise codes	- <u>endemic A09.0</u>
Add subterm	- epidemic A09.0
	- flagellate A07.9 - functional K59.1
	- full cholai K39.1 - following gastrointestinal surgery K91.8
	Poliowing gastromestrial surgery K91.8 psychogenic F45.3
	psychogenic 143.3 - Giardia lamblia A07.1
	- giardial A07.1
	- infectious A09.0
	- inflammatory A09 A09.9
Revise codes and	- infectious A09.0
add subterms	- noninfectious K52.9
	- mycotic NEC B49+ K93.8*
	- neonatal (noninfectivenoninfectious) P78.3
Revise text	- nervous F45.3
	- <u>noninfectious</u> K52.9
	— presumed noninfectious K52.9

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Delete text	- protozoal A07.9
	specified NEC A07.8
	- psychogenic F45.3 - specified
	bacterium NEC A04.8
	virus NEC A08.3
	- strongyloidiasis B78.0
	- toxic K52.1
	- trichomonal A07.8
	- tropical K90.1
	- tuberculous A18.3+ K93.0* - viral (<i>see also</i> Enteritis, viral) A08.4
Add subterm	Diarrhea, diarrheal (disease) (infantile) A09.9
Add subterm	- due to
	– bacteria A04.9
	specified NEC A04.8
	Campylobacter A04.5
	<u>– – Clostridium difficile A04.7</u>
	Clostridium perfringens (C) (F) A04.8
	 - Cryptosporidium A07.2 - Escherichia coli A04.4
	Diarrhea, diarrheal (disease) (infantile) A09.9
	Diarrilea, marrilear (misease) (initantile) A07.7
A 1.1 . 1.4	- dietetic K52.2
Add subterms	- drug-induced K52.1
	- due to
	bacteria A04.9
	specified NEC A04.8
	Campylobacter A04.5
	Clostridium perfringens (C) (F) A04.8 Cryptosporidium A07.2
	Crypiosportatum A01.2 drugs K52.1
Revise codes	Dilatation
	– eustachian tube, congenital Q17.8-Q16.4
Add dagger	Diphtheria, diphtheritic (gangrenous) (hemorrhagic) A36.9
asterisk	– neurological complication A36.8

INSTRUCTION	ALPHABETIC INDEX ENTRIES
combinations	<u>— neuritis A36.8 † G59.8*</u> — paralysis A36.8 † G63.0*
Revise code	Disease, diseased - see also Syndrome - diarrheal, infectious NEC A09 A09.0
Add subterms	Disease, diseased — see also Syndrome - arterial — see also Disease, artery - occlusive I77.1 mesenteric (chronic) K55.1 acute K55.0
Revise subterms	Disease, diseased — see also Syndrome end-stage renal kidney N18.0 N18.5
and codes:	- endocrine glands or system NEC E34.9 - endomyocardial (eosinophilic) I42.3 glomerular (see also Glomerulonephritis) N05 with edema — see Nephrosis - chronic N03 kidney (functional) (pelvis) (see also Disease, renal) N28.9
	Note: Where a term is indexed only at the three-character level, e.g. N01, reference should be made to the list of fourth-character subdivisions in Volume 1 at N00-N08. with
Add note and subterms:	edema – see Nephrosis glomerular lesion – see Glomerulonephritis with edema – see Nephrosis interstitial nephritis N12 - acute — see Nephritis, acute - chronic N18.9 end-stage N18.5 stage 1 N18.1 stage 2 N18.2 stage 3 N18.3 stage 4 N18.4 stage 5 N18.5 - cystic (congenital) Q61.9

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Delete note and subterms:	- end-stage (failure) N18.5 - fibrocystic (congenital) Q61.8 - hypertensive (see also Hypertension, kidney) I12.9 - rend-stage (failure) I12.0 - rend (fare solve) schistosomaissis (biharziasis) B65† N29.1* - rend(ly progressive N01 tubular (see also Nephritis, tubulo-interstitial) N12 - kissing B27.9 - renal (functional) (pelvis)- See Disease, kidney N28.9 Note: Where a term is indexed only at the three character level, e.g. N01., reference should be made to the list of fourth character subdivisions in Volume 1 at N00 N08. with edema - see Nephrosis glomerular lesion - see Glomerulonephritis with edema - see Nephrosis interstitial nephritis N12 - acute - see Nephrosis - interstitial nephritis N12 - acute - see Nephrosis - stage 1 N18.1 - stage 2 N18.2 - stage 3 N18.3 - stage 3 N18.3 - stage 3 N18.3 - stage 3 N18.4 - stage 5 N18.5 - stage 1 N18.4 - stage 5 N18.5 - fibrocystic (congenital) Q61.9 - end stage (failure) N18.5 - fibrocystic (congenital) Q61.8 - hypertensive (see also Hypertension, kidney) 112.9 - end stage (failure) N18.5 - rapidly progressive N01 tubular (see also Nephritis, tubulo interstitial) N12 - Disease, diseased - continued

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Revise codes and	 neoplastic (malignant), generalized (M8000/6) C80 C79.9
add subterms:	 – primary site not indicated C80.9 – primary site unknown, so stated C80.0
	Disease, diseased - see also Syndrome
	- sickle cell D57.1
Revise code:	- thalassaemia D56.8 <u>D57.2</u>
Delete	Disease, diseased – see also Syndrome
morphology codes	
and add subterms	– alpha heavy chain (M9762/3) C88.4 <u>3</u>
	– gamma heavy chain (M9763/3) C88.2
	- hand, foot and mouth B08.4
	- Hand-Schueller-Christian C96.5
	- heavy chain
	alpha (M9762/3) C88. <u>13</u>
	gamma (M9763/3) C88.2
	<u> Mu C88.2</u> NEC C88.2
	<u>— NEC C88.2</u> — immunoproliferative (malignant) (M9760/3) C88.9
	- small intestinal (M9760/3) C88.3
	- specified NEC C88.7
	- lymphoproliferative (M9970/0) D47.9
	– T–gamma (M9768/1) D47. 7 9
	- X-linked D82.3
	 myelodysplastic and myeloproliferative, not classifiable C94.6
	– myeloproliferative (chronic) (M9960/1) D47.1
	– systemic tissue mast cell (M9741/3) C96.2 <u>D47.0</u>
Add non-essential	Disease, diseased (see also Syndrome)
modifier, index	···
leadterm and	- liver (chronic)(organic) K76.9
subterms	
	 - drug-induced (idiosyncratic)(toxic)(predictable)(unpredictable) - see Disease, liver toxic - fatty, nonalcoholic K76.0
	· · · · · · · · · · · · · · · · · · ·
	inflammatory K75.9
	specified K75.8
	- nonalcoholic, fatty K76.0
Add and revise	Disease, diseased (see also Syndrome)

INSTRUCTION	ALPHABETIC INDEX ENTRIES
subterms	- kidney (functional) (pelvis) (see also Disease, renal) N28.9
	cystic (congenital) Q61.9 multicystic Q61.4
	polycystic Q61.3
	- – fibrocystic (congenital) Q61.8
Add asterisk code	Disease, diseased (see also Syndrome)
	- woolsorter's A22.1† J17.0* Disease, diseased - continued
Delete subterm	- neoplastic (malignant), generalized (M8000/6) C79.9
	— primary site not indicated C80.9 — primary site unknown, so stated C80.0
	Disease, diseased - see also Syndrome
	- kidney (functional) (pelvis) N28.9
Add code and	complicating pregnancy O26.8
subterms	with hypertension (preexisting) O10.2
	affecting fetus or newborn P00.1
	- renal (functional) (pelvis) (see Disease, kidney)
	Disease, diseased—continued
Add subterms	- silo-filler's J68.8 - bronchitis J68.0
Add subterms	- pneumonitis J68.0
	pulmonary edema J68.1
	Disease, diseased - see also Syndrome
D : 1	
Revise code	obstructive (chronic) J44.9
	with
	 emphysema J44. <u>J43.</u> -
Add subterm	Disease, diseased
	••••
	- Keshan E59
	- kidney (functional) (pelvis) N28.9 - complicating pregnancy O26.8

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	in (due to) schistosomiasis (bilharziasis) B65† N29.1* puerperal, postpartum O90.8
Add subterm	Disease, diseased — see also Syndrome
	- hepatobiliary K83.9
	- kidney (functional) (pelvis) N28.9 Note: Where a term is indexed only at the three-character level, e.g. N01, reference should be made to the list of fourth-character
	subdivisions in Volume 1 at N00-N08.
	cystic (congenital) Q61.9
Add subterm	diabetic (acute)(any stage)(chronic)(See also E10-E14 with fourth character .2) E14.2† N08.3* end-stage (failure) N18.5
Add subterm	Disorder (of)- see also Disease
	- metabolism
	phosphatases E83.3 phosphorus E83.3
Add non-essential	Disorder (of)-see also Disease
modifier:	- developmental (global delay) F89
Delete	Disorder (of) (see also Disease)
morphology code	– glomerular (in) N05.9
and apostrophe	− multiple myeloma (M9732/3) C90.0† N08.1*
1 1	– Waldenström's macroglobulinemia (M9761/3) C88.0† N08.1*
	
	- tubulo-interstitial (in)
	leukemia NEC (M98003/) C95.9† N16.1*
	 lymphoma NEC (M9590/3) C85.9† N16.1* multiple myeloma (M9732/3) C90.0† N16.1*
	Disorder (of) (see also Disease)
	– psychotic (due to) (see also Psychosis) F29
Add and revise	alcohol(- induced), alcoholic (immediate) F10.5
Add and revise	intoxication (acute) F10.5

INSTRUCTION	ALPHABETIC INDEX ENTRIES
subterms	<u>– – residual and late onset F10.7</u>
	- drug induced (immediate)(see also F11–F19 with fourth character .5)
	F19.5
	residual and late onset (<i>see also</i> F11–F19, with fourth character .7) F19.7
	<u> residual and late onset</u>
	<u> alcohol-induced F10.7</u> drug-induced (<i>see also</i> F11-F19 with fourth character .7) F19.7
	Disorder (of) – see also Disease
Modify subterms	– male
and add subterms	erectile F52.2
	organic origin NEC N48.4 psychogenic F52.2
	- hypoactive sexual desire F52.0
	Dissection
	- aorta (any part)(ruptured) I71.0
A 1.1 - 1.4	- artery <u>NEC 172.9</u>
Add subterms:	carotid I72.0 cerebral (nonruptured) I67.0
	ruptured (<i>see also</i> Hemorrhage, subarachnoid) I60.7
	iliac (ruptured) I72.3
	<u> limb (ruptured)</u>
	<u>lower I72.4</u>
	<u> upper I72.1</u> <u> precerebral NEC I72.5</u>
	acquired (ruptured) I72.5
	carotid I72.0
	<u> vertebral I72.5</u>
	congenital (nonruptured) Q28.1
	renal (ruptured) I72.2 specified (ruptured) NEC I72.8
	specified (ruptured) NEC 172.8 -traumatic (complication)(early), specified site - see Injury, blood vessel, vascular
	- vascular I99 NEC I72.9
	- wound - see Wound, open

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Revise and add subterms	Dissection - aorta (any part)(ruptured) I71.0 - artery NEC I72.9 - basilar I72.5 - carotid I72.0 - vertebral I72.6 precerebral NEC I72.5 acquired NEC (ruptured) I72.5 acquired NEC (ruptured) I72.5 carotid I72.0 vertebral I72.5
Add lead term	Distal intestinal obstruction (syndrome) E84.1 Distoma hepaticum infestation B66.3
Revise code	Distortion (congenital) – eustachian tube Q17.8 Q16.4
Modify subterm	Dropped - dead, unexplained R96.0
Revise subterm	Duchenne's - disease or syndrome - motor neuron disease G12.2 — due to or associated with — motor neuron disease G12.2 — muscular dystrophy G71.0 - nuscular dystrophy G71.0 - locomotor ataxia (syphilitic) A52.1
Revise modifiers and code	Dysentery, dysenteric (catarrhal) (diarrhea) (epidemic) (hemorrhagic) (infectious) (sporadic) (tropical) A09 A09.0 - amebic (see also Amebiasis) A06.0 - with abscess - see Abscess, amebic - acute A06.0 - chronic A06.1 - arthritis A09.9+ M01.8*M03* - bacillary A03.9+ M01.8*M03* - infectious A09.0+ M01.8*M03* - noninfectious K52.9+ M01.8* - bacillary A03.9

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	arthritis A03.9+ M01.3*
	Boyd A03.2 Flexner A03.1
	Schmitz(-Stutzer) A03.0
	- Shiga(-Kruse) A03.0
	- Shigella A03.9
	boydii A03.2
	dysenteriae A03.0
	flexneri A03.1
	group A A03.0
	group B A03.1
	group C A03.2 group D A03.3
	group D A03.3 sonnei A03.3
	specified type NEC A03.8
	- Sonne A03.3
	specified type NEC A03.8
	Dysfunction
	- sexual (due to)
	alcohol F10.8
Revise code:	opioid F11.0 <u>F11.8</u>
Modify subterms	Dysfunction
and add subterms	<u>– erectile (psychognic) F52.2</u>
	organic origin NEC N48.4
Revise code	Dysfunction - hemoglobin D75.8 D58.2
	Dysfunction
Revise code	- hypophysis E23. 3 7
	Dyslexia R48.0
	- developmental F81.0
Add lead term and	<u>Dyslipidemia E78</u>
subterms:	- depressed HDL cholesterol E78.6
	- elevated fasting triglycerides E78.1
Add subterm	Dyspepsia (allergic)(congenital)(functional)(gastrointestinal)(occupational)(reflex) K30 R10.1 - atonic K30
	- functional (allergic)(congenital)(gastrointestinal)(occupational)(reflex) K30
	- 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	- nervous F45.3
	- neurotic F45.3
	- psychogenic F45.3 Dysplasia - see also Anomaly
Revise subterm:	- metaphyseal (Jansen's)(McKusick's)(Schmid's) Q78.5
110 / 150 500 101111	Dysplasia - see also Anomaly
	- polyostotic fibrous Q78.1
Add subterms:	- prostate (low grade) N42.3
	<u> high grade D07.5</u>
	- renal Q61.4
Add and revise	Dysplasia – kidney (multicystic) Q61.4
subterms	- renal (multicystic) Q61.4
	Dystocia O66.9
	- affecting fetus or newborn P03.1
Add non-essential	- cervical (hypotonic) (failure of cervical dilatation) O62.20
modifier and	affecting fetus or newborn P03.6
revise subterms:	due to abnormality of cervix O65.5
	— primary O62.0
	— secondary O62.1
Revise code:	Eaton-Lambert syndrome_C80C80† G73.1*
Revise cross	Eaton-Lambert syndrome (see also Neoplasm) C80† G73.1*
reference	- unassociated with neoplasm G70.8
	Eclampsia, eclamptic (coma) (convulsions) (delirium) (with pre existing or pregnancy related hypertension) NEC O15.9
Delete subterm	- male R56.8 - not associated with pregnancy or childbirth R56.8
	- pregnancy O15.0
	- affecting fetus or newborn P00.0
	Ecthyma L08.0
Revise code:	– gangrenosum L88 L08.0
Add subterm	Ectropion H02.1
	– cervix N86
	– with cervicitis N72
	congenital Q51.8
Add dagger	Egyptian splenomegaly B65.1 † D77*

INSTRUCTION	ALPHABETIC INDEX ENTRIES
symbol and asterisk to code at lead term	
Add subterms	Elevated, elevation - venous pressure I87.8 - white blood cells NEC R72
Revise subterms:	Embolism (septic) I74.9 - pyemic (multiple) (see also Septicemia Sepsis) A41.9 - septicemic — see Septicemia Sepsis
Add subterms	Encephalitis (chronic) (hemorrhagic) (idiopathic) (nonepidemic) (spurious) (subacute) G04.9 - in (due to) - cytomegalovirus B25.8† G05.1* - enterovirus A85.0† G05.1* - herpes (simplex) virus B00.4† G05.1*
Add subterm	human metapneumovirus A85.8† G05.1* Encephalitis - continuedRasmussen (subacute) G04.8
Revise subterms	Encephalitis - in (due to) - influenza (specific virus not identified) (J09†, J10.8†, J11.8†) G05.1* - avian influenza virus identified J09† G05.1* - other influenza virus identified J10.8† G05.1*
Revise subterms	Encephalopathy (acute) - in (due to) - influenza (specific virus not identified) (J09†, J10.8 J11.8†) G94.8* avian influenza virus identified J09† G94.8* other influenza virus identified J10.8† G94.8*
Delete index entry as follows	Encephalopathy – postcontusional F07.2 – current injury \$\frac{506.2}{2}\$—see Contusion, brain
Add dagger symbol	Encephalopathy (acute) G93.4 - in (due to) - influenza (J09†, J10.8†, J11.8†) G94.8*
Add subterm	Endophthalmitis (acute) (infective) (metastatic) (subacute) H44.0 — with associated postprocedural bleb H59.8

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Add subterm	Entanglement
	– umbilical cord(s) O69.2
	with compression O69.2
	– affecting fetus or newborn P02.5
	– around neck (with compression) O69.1
	<u>– – without compression O69.8</u>
	− − of twins in monoamniotic sac O69.2
Revise	Enteritis (acute) (diarrheal) (epidemic) (hemorrhagic) (presumed infectious) (septic) (see also Note at category A09) A09 A09.9
	-with septicemia NEC A09.0
	<u>- acute A09.9</u>
Add subterms	<u> bloody A09.0</u>
	hemorrhagic A09.0
	- allergic K52.2
	- amebic (see also Amebiasis) A06.0
	with abscess - see Abscess, amebic
	nondysenteric A06.2
	chronic A06.1
	with abscess - see Abscess, amebic
	nondysenteric A06.2
	nondysenteric A06.2
	- bacillary NEC A03.9
	- bacterial A04.9
	specified NEC A04.8
	- candidal B37.8
	- Chilomastix A07.8
	- chronic (noninfectivenoninfectious) K52.9
Revise	ulcerative K51.9
	- coccidial A07.3
	- dietetic K52.2
	- due to
	food hypersensitivity K52.2
	infectious organism (bacterial) (viral) - see Enteritis, infectious
	Yersinia enterocolitica A04.6
	- eltor A00.1
	- <u>epidemic A09.0</u>
Add subterm and	- gangrenous (see also Enteritis, infectious) A09.0
revise codes	- giardial A07.1

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	- infectious NEC A09.0
	due to
	adenovirus A08.2
	Aerobacter aerogenes A04.8
	Arizona (bacillus) A02.0
	bacteria NEC A04.9
	specified NEC A04.8
	Campylobacter A04.5
	Clostridium perfringens A04.8
	Enterobacter aerogenes A04.8
	enterovirus A08.3
	Escherichia coli A04.4
	enteroaggregative A04.4
	enterohemorrhagic A04.3
	enteroinvasive A04.2
	enteropathogenic A04.0
	enterotoxigenic A04.1
	specified NEC A04.4
	specified
	bacteria NEC A04.8
	virus NEC A08.3
	Staphylococcus A04.8
	virus NEC A08.4
	specified type NEC A08.3
	Yersinia enterocolitica A04.6
	specified organism NEC A08.5
	- influenzal (specific virus not identified) J11.8
	influenza virus identified J10.8
	- ischemic K55.9
	acute K55.0
	chronic K55.1
	- microsporidial A07.8
	- mucomembranous, myxomembranous - <i>see</i> Syndrome, irritable bowel
	- mucous - <i>see</i> Syndrome, irritable bowel
	- necroticans A05.2
	- necrotizing of fetus or newborn P77
	- neurogenic - see Syndrome, irritable bowel

INSTRUCTION	ALPHABETIC INDEX ENTRIES
D • • • • • • • • • • • • • • • • • • •	- noninfectious K52.9
Revise subterm	- parasitic NEC B82.9
	- paratyphoid (fever) (<i>see also</i> Fever, paratyphoid) A01.4 - presumed noninfectious K52.9
Delete text	- presumed nonninectious K32.9 - protozoal A07.9
Delete text	- specified NEC A07.8
	- regional (of) K50.9
	- intestine K50.9
	large (colon or rectum) K50.1
	with small intestine K50.8
	small (duodenum, ileum or jejunum) K50.0
	with large intestine K50.8
	- rotaviral A08.0
	- Salmonella, salmonellosis (arizonae) (cholerae-suis) (enteritidis) (typhimurium) A02.0
	- segmental K50.9
	- <u>septic A09.0</u>
	- Shigella (see also Infection, Shigella) A03.9
Add subterm	- spasmodic, spastic - <i>see</i> Syndrome, irritable bowel
	- staphylococcal A04.8
	- toxic K52.1
	- trichomonal A07.8
	- tuberculous (see also Tuberculosis) A18.3+ K93.0* - typhosa A01.0
	- typhosa A01.0 - ulcerative (chronic) K51.9
	- specified NEC K51.8
	- viral A08.4
	adenovirus A08.2
	enterovirus A08.3
	Rotavirus A08.0
	small round structured A08.1
	virus specified NEC A08.3
	Enteritis (diarrheal)(hemorrhagic) A09.9
Daviga guhtarras	– influenzal (specific virus not identified) J11.8
Revise subterms	– – avian <u>certain identified</u> influenza virus identified J09
	 – other influenza virus identified J10.8
Delete index entry	Enteritis (diarrheal) (hemorrhagic) A09.9
	-with septicemia NEC A09.0

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	Enteritis (diarrheal) (hemorrhagic) A09.9
Add subterms	 - dietetic K52.2
Add subterilis	- drug-induced K52.1
	- due to drugs K52.1
	food hypersensitivity K52.2
Add subterms	Enteritis (diarrheal) (hemorrhagic) A09.9
	infectious NEC A09.0
	due to
	Campylobacter A04.5 Clostridium difficile A04.7
	Clostridium perfringens A04.8
	- staphlococcal A04.8 - toxic <u>NEC</u> K52.1
	due to Clostridium difficile A04.7
Revise code	Enterocolitis (see also Enteritis) A09 A09.9
	- due to <i>Clostridium difficile</i> A04.7 - fulminant ischemic K55.0
	- hemorrhagic (acute) K55.0
Add subterm	- infectious NEC A09.0
	- ischemic K55.9
	- necrotizing due to <i>Clostridium difficile</i> A04.7
	fetus or newborn P77
Add subterm	- noninfectious NEC K52.9
	- ulcerative (chronic) K51.0
Add subterm	Eosinophilia (allergic) (hereditary) D72.1 - peritoneal – see Peritonitis
	- perionear – see Fernolius - pulmonary NEC J82
	- tropical (pulmonary) J82
Delete	Erythremia (acute) (M9841/3) C94.0
	– chronic (M9842/3) C94.1 <u>D45</u>

INSTRUCTION	ALPHABETIC INDEX ENTRIES
morphology codes and revise code	- secondary D75.1
Delete morphology code	Erythrocythemia (M9841/3) – see Erythremia
Delete morphology code	Erythroleukemia (M9840/3) C94.0
Add leadterm, subterm and codes	Fahr-Volhard disease (of kidney) I12.9 - with renal insufficiency I12.0
Add leadterm and code	Fahr's disease (of brain) G23.8
Add subterms Add subterms	Failure, failed – continued heart (acute) (sudden) I50.9 - with acute pulmonary oedema — see Failure, ventricular, left decompensation (see also Failure, heart, congestive) I50.9 dilatation — see Disease, heart other organ failure, code each site (see also Failure organ multiple NEC) - hepatic K72.9 - with other organ failure, code each site (see also Failure organ multiple NEC) - acute or subacute K72.0 - myocardial, myocardium (see also Failure, heart) I50.9 - organ — see Failure, by site - multiple NEC R68.8 - orgasm (female)(male)(psychogenic) F52.3 - renal N19 - with diabetes — see Diabetes, with, failure, renal - hypertension (see also Hypertension, kidney) I12.0 hypertensive heart disease (conditions in I11) I13.1 with heart failure (congestive) I13.2 other organ failure, code each site (see also Failure organ multiple NEC)

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	- respiration, respiratory J96.9
	with
	other organ failure, code each site (see also Failure organ multiple NEC)
	acute J96.0
4.11 1.	centre G93.8
Add subterms	Failure, failed
	- heart (acute) (sudden) I50.9
	- Heart (acute) (sudden) 150.9
	hypertensive (see also Hypertension, heart) I11.0
	with renal disease I13.0
	with renal failure I13.2
	<u> ischemic I25.5</u>
	left (ventricular) (see also Failure, ventricular, left) I50.1
Add non-essential	Failure, failedcontinued
modifier and	- heart (acute) (sudden) (senile) I50.9
delete subterm:	senile R54
	Failure, failed
Delete cross	- kidney — see Failure, renal N19
reference and add	<u> with</u>
subterms:	hypertension (see also Hypertension, kidney) I12.0
	hypertensive
	heart disease (conditions in II1) II3.1
	with heart failure (congestive) I13.2 kidney disease (see also Hypertension, kidney) I12.0
	tubular necrosis (acute) N17.0
	acute N17.9
	with
	cortical necrosis N17.1
	medullary necrosis N17.2
	tubular necrosis N17.0
	following labour and delivery O90.4
	specified NEC N17.8
	<u> chronic N18.9</u>
	<u> end-stage N18.5</u>
	hypertensive (see also Hypertension, kidney) I12.0
	<u> stage 1 N18.1</u>

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Delete code, add cross reference and delete subterms:	stage 2 N18.2stage 3 N18.3stage 4 N18.4stage 5 N18.5stage 5 N18.5compenital P96.0end-stage (chronic) N18.5followingabortion (subsequent episode) O08.4current episode — see Abortioncrushing T79.5ctopic or molar pregnancy O08.4labour and delivery (acute) O90.4bypertensive (see also Hypertension, kidney) I12.0postprocedural N99.0renal N49 — see Failure, kidney

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	— labor and delivery (acute) O90.4
	— hypertensive (see also Hypertension, kidney) 112.0 — postprocedural N99.0
	- trial of labor (with subsequent caesarean section) O66.4
Revise cross-	affecting fetus or newborn P03.1
reference:	- urinary — see Failure, renal <u>kidney</u>
	Failure, failed
Delete subterm	- kidney N19
	with
Add subterm	Failure, failed
Add subterin	ranure, raneu
	- kidney N19
	congenital P96.0
	diabetic (acute) (any stage)(chronic)(See also E10-E14 with fourth character .2) E14.2† N08.3*
	end-stage (chronic) N18.5
Modify subterm	Fatal syncope, unexplained R96.0
Add non-essential	Fatty – see also condition
modifier	– liver (nonalcoholic) NEC K76.0
	alcoholic K70.0
	Fever R50.9
Revise cross	- putrid — see Septicaemia Sepsis
reference:	- pyemic — see Septicaemia <u>Sepsis</u>
G 111	- septic — see Septicaemia-Sepsis Fever
Correct spelling	rever - remittent (congestive) (see also Ma r -laria) B54
in cross reference	
Add dagger	Fever - typhoid (abortive) (hemorrhagic) (intermittent) (malignant) A01.0
asterisk codes	- typhoid (abortive) (hemormagic) (intermittent) (manghant) A01.0
	gastro-intestinal perforation A01.0† K93.8*
	with peritonitis A01.0† K67.8*
	pneumopathy A01.0† J17.0*
Revise code at	Fever R50.9

ALPHABETIC INDEX ENTRIES
- Mediterranean <u>A23.0-A23.9</u>
Fibrillation
- atrial or auricular (established) I48 <u>.9</u>
<u> chronic I48.2</u>
paroxysmal I48.0 persistent I48.1
- persistent 146.1 - cardiac I49.8
Fibroadenoma (M9010/0)
- prostate N40 <u>D29.1</u>
Fibroma (M8810/0) — see also Neoplasm, connective tissue, benign
- prostate N40 <u>D29.1</u>
Fibrosis, fibrotic
- cystic (of pancreas) E84.9
 – with – combined manifestations E84.8
distal intestinal obstruction (syndrome) E84.1
intestinal manifestations E84.1
Fibrosis, fibrotic
••••
- cystic (of pancreas) E84.9
with
combined manifestations E84.8
 distal intestional obstruction (syndrome) E84.1 intestinal manifestations E84.1
pulmonary manifestations E84.0
- specified NEC E84.8
Fibrosis, fibrotic
- lung (atrophic) (capillary) (chronic) (confluent) (massive) (perialveolar) (peribronchial) J84.1
with
asbestosis J61
<u> amianthosis J61</u>
Fistula L98.8
- arteriovenous (acquired) (nonruptured) I77.0 coronary I25.4
congenital O24.5

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	- congenital, site not listed - see Anomaly, by site
	- coronary, arteriovenous I25.4 congenital Q24.5
Add subterms	Fistula L98.8
Add subternis	- bronchocutaneous, bronchomediastinal, bronchopleural, bronchopleuromediastinal
	(infective) J86.0
	– – tuberculous NEC A16.4
	<u> with bacteriological and histological confirmation A 15.5</u>
	– pleura, pleural, pleurocutaneous or pleuroperitoneal J86.0
	 - tuberculous NEC A16.5 - with bacteriological and histological confirmation A15.6
A 11 14	FitzhughFitz Hugh and Curtis syndrome (gonoccal)A54.8† K67.1*
Amend lead term and add subterms	- chlamydial A74.8† K67.0*
and add subterms	- gonococcal A54.8† K67.1*
Add subterms,	Flutter
modify codes and	- atrial or auricular I48.9
delete subterm	- heart I49.8
	atrial I48 <u>.9</u>
	atypical I48.4
	<u> type I I48.3</u> type II I48.4
	typical I48.3
	- ventricular I49.0
	-ventricular I49.0
Add optional	Forestier's disease M48.1 (rhizomelic pseudopolyarthritis) M35.3
modifiers,	- ankylosing hyperostosis M48.1
subterm and code:	
Revise code	Fracture (abduction)(adduction)(avulsion)(comminuted)(compression) (dislocation)(oblique)(separation) T14.2 - march \$\frac{892.3}{2} \frac{M84.3}{2}\$
	Fracture (abduction) (adduction) (avulsion) (comminuted) (compression) (dislocation) (oblique) (separation) T14.2
	- tibia, tibial (shaft) (with fibula) S82.2
Revise and add	condyles S82.1 distal end S82.3
subterm	distal end \$82.5
	lower S82.3
	upper S82.1

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	head (involving knee joint) S82.1 intercondyloid eminence S82.1
	involving ankle or malleolus S82.5 lower end (with fibula) S82.3
	<u> plateau S82.1</u>
	proximal end S82.1 tuberosity S82.1
	upper end or extremity S82.1
Add subterms	Fracture (abduction) (adduction) (avulsion) (comminuted) (compression) (dislocation) (oblique) (separation) T14.2
	- base of skull S02.1
	- <u>basicervical (basal) (femoral) S72.0</u>
	- femur, femoral S72.9
	<u>basicervical (basal) S72.0</u>
	birth injury P13.2 condyles, epicondyles S72.4
Delete	Franklin's disease (M9763/3) C88.2
morphology code	Gammopathy
Delete morphology code	- associated with lymphoplasmacytic dyscrasia (M9765/1) D47.2
and add non-	– monoclonal (of undetermined significance) (M97651/1)D47.2
essential modifier	— of undetermined significance D89.2
Delete last	
subterm	Consume consumers (Aur.) (maint) (alim) (alean) (and alim) (alean) (and alim) (alim) (alean) (and alim)
	Gangrene, gangrenous (dry) (moist) (skin) (ulcer) (see also Necrosis) R02
	- appendix K35.9 <u>K35.8</u>
	with
	— perforation or rupture- K35.0
	peritoneal abscess K35.1 <u>K35.3</u> peritonitis, generalized K35.2
	with mention of perforation or rupture K35.2
Revise subterms:	peritonitis, localized K35.9 K35.3
	with mention of perforation or rupture K35.0 <u>K35.3</u>
	generalized K35.0

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Revise subterms	Gangrene, gangrenous (dry)(moist)(skin)(ulcer)(see also Necrosis) R02
	Was a
	- appendix K35.8
	with
	peritoneal abscess K35.3 peritonitis , generalized NEC K35.2 K35.3
	— with mention of perforation or rupture K35.2
	— peritonitis, localized K35.3
	- with mention of perforation or rupture K35.3
	generalized (with perforation or rupture) K35.2
	localized (with perforation or rupture) K35.3
Revise	Gastroenteritis (acute) (epidemic) (presumed infectious) (see also Note at category A09) A09 (see also Enteritis) A09.9
	- with septicemia NEC A09.0
	- allergic K52.2
Revise text	- chronic (noninfectivenoninfectious) K52.9
	- dietetic K52.2
	- due to
	Cryptosporidium A07.2
	food poisoning (<i>see also</i> Intoxication, foodborne) A05.9 radiation K52.0
	- eosinophilic K52.8
Add subterm	- epidemic A09.0
Tida sasterini	- food hypersensitivity K52.2
	- infectious (see also Enteritis, infectious) A09.0
	- nonbacterial, of infancy A08.5
Add subterm	- noninfectious K52.9
	specified NEC K52.8
	- rotaviral A08.0
	- salmonella A02.0
Add subterm	- septic A09.0
	- toxic K52.1
	- viral A08.4
	acute infectious A08.3
	epidemic A08.1 type Norwalk A08.1
	type Norwalk A08.1 infantile (acute) A08.3
	Norwalk agent A08.1
	1101 Walk agon 1100.1

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	rotaviral A08.0 severe, of infants A08.3 specified type NEC A08.3
Delete subterm	Gastroenteritis (acute) (see also Enteritis) A09.9 -with septicemia NEC A09.0
Add subterms	Gastroenteritis (acute) (see also Enteritis) A09.9 - with septicemia NEC A09.0 - allergic K52.2 - chronic (noninfectious) K52.9 - dietetic K52.2 - drug-induced K52.1 - due to - Cryptosporidium A07.2 - drugs K52.1 - food poisoning (see also Intoxication, foodborne) A05.9
Revise code	Gastroenteropathy (see also Gastroenteritis) A09 A09.9
Add subterms	- acute, due to Norwalk agent A08.1 - infectious A09.0 - noninfectious K52.9
Revise	Gastrojejunitis, noninfective noninfectious (presumed) (see also Enteritis and note at category A09) K52.9
Delete morphology codes and revise codes	Germinoblastoma (diffuse) (M9676/3) C83.2 C85.9 – follicular (M9692/3) C82.49
Revise subterms:	Glomerulonephritis (see also Nephritis) N05 in (due to) schistosomiasis B65† N08.0* septicemia sepsis A41† N08.0*
Add dagger/asterisk combination:	Glomerulonephritis (see also Nephritis) N05 - in (due to) - septicemia A41 N08.0* streptococcal A40† N08.0*
Delete morphology codes and apostrophe	Glomerulonephritis (see also Nephritis) N04.– – in (due to) – multiple myeloma (M9732/3) C90.0† N08.1* – Waldenström²s macroglobulinemia (M9761/3) C88.0† N08.1*
Revise subterm	Glomerulonephritis (see also Nephritis) N05

INSTRUCTION	ALPHABETIC INDEX ENTRIES
and code	- sclerosing, diffuse (see also Failure, kidney, chronic) N19N18.9
Revise cross reference and add	Glottitis – see Glossitis (see also Laryngitis) J04.0
code:	
Revise code	Glucagonoma-(M8152/0)
	- pancreas D13.7
	<u> benign (M8152/0) D13.7</u>
	uncertain or unknown behavior (M8152/1) D37.7
	malignant (M8152/3) C25.4
	- specified site NEC— see Neoplasm, benign
	benign (M8152/0) - see Neoplasm, benign
	uncertain or unknown behavior (M8152/1) - see Neoplasm, uncertain or unknown behavior malignant (M8152/3) - see Neoplasm, malignant
	- mangnant (M8132/3) – see Neopiashi, mangnant - unspecified site- <u>D13.7</u>
	- unspectified site_ D13.7 - benign (M8152/0) D13.7
	- uncertain or unknown behavior (M8152/1) D37.7
	- malignant (M8152/3) C25.4
	Gout, gouty M10.9
	– syphilitic A52.7† M14.8*
Revise codes	– tophi NEC <u>M10.9</u> <u>M10.0</u>
Revise codes	− − ear M10.9 <u>M10.0</u> † H62.8*
Delete	Granuloma L92.9
morphology	- bone M86.8
codes, revise	eosinophilic D76.0 <u>C96.6</u>
codes and	from residual foreign body M86.8
subterms	<u>lung C96.6</u>
Delete subterm	 eosinophilic D76.0 <u>C96.6</u> – bone D76.0 <u>C96.6</u>
	bone D76.0 <u>C96.6</u> lung D76.0 C96.6
	· —
	 – Hodgkin 's (M9661/3) C81. 7 9
	- lung (infectious) (see also Fibrosis, lung) J84.1
	- coccidioidal B38.1† J99.8*
	– eosinophilic D76.0 <u>C96.6</u>
Delete	Granulomatosis L92.9
morphology code,	– lymphoid (M9766/1) D47.7 <u>C83.8</u>

INSTRUCTION	ALPHABETIC INDEX ENTRIES
add subterm and	<u>– lymphomatoid C83.8</u>
revise code Revise code and	Gumboil K04.67
add subterm	- with sinus K04.6
Revise code	Haff disease T56.1 T61.2
Revise code in	Hallux
subterm	– valgus (acquired) M20.1
	– – congenital Q66. <u>86</u>
	– varus (acquired) M20.3
	congenital Q66.3
Revise code	Hand–Schüller–Christian disease or syndrome D76.0 C96.5
Add subterm and	Health
code	- services provided because (of)
	reduced mobility Z74.0
Revise code and	Heilmeyer–Schoner disease (M9842/3) C94.1 D45
delete	
morphology code	Helicobacter pylori, as cause of disease classified elsewhere B98.0
Add lead term	
Add non-essential	HELLP syndrome (hemolysis, elevated liver enzymes and low platelet count) O14.42
modifiers and	
revise code	Hematoma (traumatic) (skin surface intact) (see also Injury, superficial) T14.0
	- penis (nontraumatic) N48.8
Revise code at	- birth injury P15.5
subterm	- perianal (nontraumatic) 184.3K64.5
	- perineal \$30.2
Add lead term	Hemicolitis, left K51.5
Add non-essential	Hemochromatosis (diabetic)(hereditary)(liver)(myocardium)(primary idiopathic)(secondary) E83.1
modifiers and	- with refractory anemia (related to alkylating agnet)(related to
delete	Epipodophyllotoxin)(related to therapy) NEC (M9982/1) D46.1
morphology code	
	Hemoglobinopathy (mixed) NEC D58.2
	- with thalassemia D56.9
Davisa anda	- sickle-cell D57.1
Revise code:	with thalassemia D56.8 D57.2

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Revise non-	Hemoglobinuria R82.3
essential	- with anemia, hemolytic, acquired (chronic) NEC D59.6
modifiers and add	– cold <u>agglutinin (disease)(agglutinin) (paroxysmal) (with Raynaud's</u>
subterm	syndrome) D59.6 D59.1
	_ – paroxysmal (with Raynaud's syndrome) D59.6
Add non-essential	Hemorrhage, hemorrhagic R58
modifiers and	- complicating
revise subterms	delivery O67.9
	affecting fetus or newborn P02.1
	associated with coagulation defect (afibrinogenemia) (hyperfibrinolysis) (hypofibrinogenemia) O67.0
	– afibrinogenemia O67.0
	affecting fetus or newborn P03.8
Revise code at	Hemorrhage, hemorrhagic R58
subterm level	- hypopharyngeal (throat) R58R04.1
Revise lead term	Hemorrhoids (bleeding) (without mention of degree) K64.9 I84.9
and subterms;	- 1st degree (grade/stage I) (without prolapse) K64.0
Add subterms.	- 2nd degree (grade/stage II) (with prolapse but retracts spontaneously) K64.1
	- 3rd degree (grade/stage III) (with prolapse that requires manual replacement) K64.2
	- 4th degree (grade/stage IV) (with prolapse that cannot be manually replaced) K64.3
	—bleeding, prolapsed, strangulated or ulcerated NEC 184.8
	- complicating
	pregnancy O22.4
	puerperium O87.2
	-external 184.5
	— bleeding, prolapsed, strangulated or ulcerated I84.4
	— thrombosed I84.3
	-internal I84.2
	— bleeding, prolapsed, strangulated or ulcerated I84.1
	— thrombosed I84.0
	- specified NEC K64.8
	- strangulated (see also Hemorrhoids, by degree) K64.8
	- thrombosed (see also Hemorrhoids, by degree) K64.8 NEC 184.7
	- ulcerated (see also Hemorrhoids, by degree) K64.8
Add subterm	Hemothorax (bacterial) (nontuberculous) J94.2
	- tuberculous NEC A16.5
	with bacteriological and histological confirmation A15.6
	Hepatitis K75.9

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Modify code	- acute NEC K72.0 <u>B17.9</u>
Add subterms	<u>- with hepatic failure (see also Failure, hepatic) K72.9</u>
	<u>– alcoholic K70.1</u>
	- infectious B15.9
	 with hepatic coma B15.0 - viral (unspecified) B17.9
	- infectious, infective (acute)(chronic)(subacute) B15.9
Modify subterms	- acute (subacute) B15.9
	chronic B18.9
	with hepatic coma B15.0
	– post-transfusion (see also Hepatitis, viral, by type type B) B16.9-B19
	– viral, virus B19.9
	- with hepatic coma B19.0
	<u>– acute (unspecified) B17.9</u>
Add subterms	- chronic B18.9
	specified NEC B18.8 type
	type $$ B B18.1
	with delta-agent B18.0
	C B18.2
	- complicating pregnancy, childbirth or puerperium O98.4
	congenital P35.3
	– – coxsackie B33.8† K77.0 □
	cytomegalic inclusion B25.1† K77.0 □
	non-A, non-B B17.8
	specified type NEC (with or without coma) B17.8
	type A B15.9
	With hepatic coma B15.0
	B B16.9
	with
	delta-agent (coinfection) (without hepatic coma) B16.1
	with hepatic coma B16.0
	hepatic coma (without delta-agent coinfection) B16.2
	C B18.2
	acute or a stated duration of less than six months B17.1
	chronic (stated duration of six months and more) B18.2

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Add subterms	E B17.2
	non-A, non-B B17.8
	<u> </u>
	<u> with hepatic coma B19.0</u>
	without hepatic coma B19.9
	 unspecified with hepatic coma B19.0
	with nepatic coma B19.0 without hepatic coma B19.9
	<u> without nepatic coma B19.9</u> acute B17.9
Add non–essential	Hepatitis K75.9
modifiers	
mounters	– fetus or newborn (giant cell) (idiopathic) P59.2
	- neonatal (giant cell) (idiopathic) (toxic) P59.2
Delete modifier	Hepatitis K75.9
Delete modifier	- anicteric , acute (viral) - see Hepatitis, viral
Revise code at	Hepatitis K75.9
subterm	- lupoid NEC K73.2 <u>K75.4</u>
Delete subterm	Hepatitis K75.9
	- acute NEC B17.9
	— with hepatic failure (see also Failure, hepatic) K72.9
	– alcoholic K70.1– infectious B17.9
	infectious B17.9 with hepatic coma B15.0
	with hepatic coma B13.0 non-viral K72.0
	- viral (unspecified) B17.9
Add subterms	vital (unspecifica) B17.5
rida suoterinis	- non-viral NEC K75.9
	acute K72.0
	- viral, virus (acute) B19.9
Revise cross-	Hernia, hernial (acquired) (recurrent) K46.9
reference	- Hesselbach's— see Hernia, femoral , by type
	Hernia, hernial (acquired)(recurrent) K46.9
Add subterm	
	- hiatal (esophageal) (sliding) K44.9

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	with
	gangrene (and obstruction) K44.1 obstruction K44.0
	congenital Q40.1
	- hypogastric — see Hernia, ventral
	- incarcerated — see also Hernia, by site, with obstruction
	with gangrene — <i>see</i> Hernia, by site, with gangrene - incisional — <i>see</i> Hernia, ventral NEC K43.2
	- metsional - see Herma, ventrar <u>IVEC K+3.2</u> with
	gangrene (and obstruction) K43.1
	<u> obstruction K43.0</u>
	- indirect (inguinal) — see Hernia, inguinal
	- mesenteric (internal) — <i>see</i> Hernia, abdomen
	- midline — see Hernia, ventral
	- muscle (sheath) M62.8
	- paraesophageal — see also Hernia, diaphragm K44.9 - congenital Q40.1
	congenital Q40.1 - parastomal K43.5
	with
	gangrene (and obstruction) K43.4
	<u> obstruction K43.3</u>
	- paraumbilical — see Hernia, umbilicus
	strangulated (see also Hernia, by site, with obstruction) K46.0
	- with gangrene (see also Hernia, by site, with gangrene) K46.1
	- subxiphoid — see Hernia, ventral
	- supra-umbilicus — see Hernia, ventral
	ventral NEC K43.9
	- with
	gangrene (and obstruction) K43.1 K43.7
	obstruction <u>K43.</u> 0- <u>K43.6</u>
A 11	Herpes, herpetic B00.9
Add non-essential modifier and	- conjunctivitis (simplex) B00.5† H13.1* - zoster B02.3† H13.1*
mounter and	ZUSICI DUZ.3 1113.1 ·

INSTRUCTION	ALPHABETIC INDEX ENTRIES
codes, and add	
subterms:	- simplex B00.9
	complicated NEC B00.8
	congenital P35.2 conjunctivitis B00.5† H13.1*
	conjunctivitis B00.5+ H15.1* external ear B00.1† H62.1*
	- zoster (see also condition) B02.9
	auricularis B02.2† H94.0*
	complicated NEC B02.8
	conjunctivitis B02.3† H13.1*
	disseminated B02.7
	Herpes, herpetic B00.9
Revise subterm:	- scrotum A60.0† N51.8*
Revise subteriii:	- septicemia sepsis B00.7 Herpes, herpetic B00.9
Add asterisk code	- whitlow B00.8† <u>L99.8*</u>
Revise cross- reference	Hesselbach's hernia— see Hernia, femoral, specified site NEC
Delete	Histiocytosis D76.3
morphology	- acute differentiated progressive (M9722/3) C96.0
codes, add	- Langerhans' cell NEC D76.0 C96.6
subterms and	<u>— multifocal (X)</u> —— multisystemic (disseminated) C96.0
revise codes	unisystemic C96.5
	- unifocal (X) C96.6
	- malignant (M9720/3) C96.18
	- mononuclear phagocytes NEC D76.1
	Langerhans' cells D76.0 C96.6
	- sinus, with massive lymphadenopathy D76.3
	- syndrome NEC D76.3
	-X NEC D76.0 C96.6
	acute (progressive) (M9722/3) C96.0 chronic D76.0 C96.6
	== cirolic B76.0 <u>C96.0</u> == multifocal <u>C96.5</u>
	- unifocal C96.6
	- multisystemic C96.0

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Revise lead term, delete morphology codes and subterms, add non-essential modifiers and subterms and revise codes	Hodgkin's - disease (M9650/3) C81.9 - lymphocytic - depletion (diffuse fibrosis) (reticular) (M9653/3) C81.3 - diffuse fibrosis (M9654/3) C81.3 - reticular (M9655/3) C81.3 - predominance (diffuse) (nodular) (M9657/3) C81.0 diffuse (M9658/3) C81.0 - nodular (M9659/3) C81.0 - lymphocytic-histiocytic predominance (M9657/3) C81.0 - mixed cellularity (M9652/3) C81.1 - nodular sclerosis (M9666/3) C81.1 - cellular phase (M9664/3) C81.1 - syncytial variant (M9667/3) C81.1 - syncytial variant (M9667/3) C81.1 - granuloma (M9661/3) C81.79 - lymphocyte depleted C81.3 - depletion (diffuse fibrosis) (reticular) C81.3 - lymphocyte-rich C81.4 - mixed cellularity C81.2 - nodular sclerosis (cellular phase) (Grade II) (Grade II) C81.1 - type not specified C81.7 - lymphocyte predominant, nodular C81.0 - paragranuloma (nodular) (M9660/3) C81.70 - sarcoma (M9662/3) C81.73
Delete indent	Hodgkin - disease C81.9 - nodular sclerosis C81.1 lymphocytic mixed cellularity C81.1
Delete apostrophe	Human - immunodeficiency virus (HIV) disease (infection) B24 resulting in Burkitt's lymphoma B21.1
	lymphoma (malignant) B21.2

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	Burkitt 's B21.1
	– – non–Hodgkin 's lymphoma NEC B21.1
Add subterm	Human – immunodeficiency virus (HIV) disease (infection) B24
	- Infinitiodeficiency virus (FITV) disease (infection) B24 - complicating pregnancy, childbirth or the puerperium O98.7
	contact Z20.6
	dementia B22.0† F02.4*
	exposure to Z20.6
	Human
	wasting syndrome B22.2
Add subterms	 metapneumovirus B97.8 resulting in
	bronchiolitis J21.1
	bronchitis J20.8
	encephalitis A85.8† G05.1*
	_ — pneumonia J12.3
	- T-cell llylmphotropic virus type-1 (HTLV-I) infection B33.3
	Hydrocele (spermatic cord) (testis) (tunica vaginalis) N43.3
A 11 17	- canal of Nuck N94.8
Add subterms	- communicating N43.2 - congenital P83.5
	- congenital P83.5
	Hypergammaglobulinemia D89.2
Delete apostrophe	- Waldenström 's D89.0
Add subterm	Hyperparathyroidism E21.3
	- specified NEC E21.2
	<u>- tertiary E21.2</u>
Add lead term and	Hyperhomocysteinemia E72.1
code	
	Hypertension, hypertensive (accelerated) (benign) (essential) (idiopathic) (malignant) (primary) (systemic) I10
	- complicating pregnancy, childbirth or puerperium O16
Delete and revise	with
subterm	— albuminuria (and oedema) (see also Pre eclampsia) O14.9
	— severe O14.1
	heart disease O10.1

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	and renal disease, pre-existing O10.3 oedema (see also Pre-eclampsia) O14.9
	severe O14.1
	renal disease, pre-existing O10.2
	affecting fetus or newborn P00.0
	essential (benign), pre-existing O10.0 with superimposed proteinuria- pre-eclampsia O11
	malignant, pre-existing O10.0
	malignant, pre-existing O10.0
	with superimposed proteinuria pre-eclampsia O11 pre-existing O10.9
	pre-existing 010.9 with superimposed proteinuria pre-eclampsia 011
	malignant O10.0
	pregnancy-induced (see also Hypertension, gestational) O13
	secondary to renal disease, pre-existing O10.4 specified NEC O10.4
	transient — see Hypertension, gestational
	- gestational (pregnancy-induced) (without significant proteinuria) O13
	— with significant proteinuria or albuminuria (and oedema) (see also Pre eclampsia) O14.9 - Goldblatt's I70.1
	Hypertension, hypertensive (accelerated) (essential) (idiopathic) (primary) (systemic) I10
Delete subterm	 with - heart involvement (conditions in I50, I51.4-I51.9 due to hypertension) (see also Hypertension, heart) I11
Belete sucterm	kidney involvement (conditions in N00-N07, N18, N19 and N26 due to hypertension) (see also Hypertension, kidney) I12
	— renal sclerosis(condition in N26) (seel also Hypertension, kidney) I12
Revise code	Hypertrophy, hypertrophic – nasal J34.8
	- nasai 334.6
	– mucous membrane (septum) J34.38
	sinus J34.8
	turbinate J34.3 Hypoplasia, hypoplastic
	- carpus Q71.8
Add subterm:	- cartilage-hair - see Dysplasia, metaphyseal
Revise codes	Hypoplasia, hypoplastic
	···

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	esophagus (congenital) Q39.8eustachian tube Q17.8-Q16.4
A 111 1.	Idiot, idiocy(congenital) F73
Add lead term	- amaurotic (Bielschowsky(-Jansky)) (familial) (infantile (late)) (juvenile (late)) (Vogt-Spielmeyer) E75.4
	IIAC (idiopathic infantile arterial calcification) Q28.8
	Heitis (see also Enteritis) A09.9
Revise code and	Ileitis (see also Enteritis) A09 A09.9
add subterms	- backwash K51.0
	- <u>infectious A09.0</u>
	- noninfectious K52.9
	- regional or terminal (ulcerative) K50.0
Revise code and	Ileocolitis (see also Enteritis) A09 A09.9
add subterms	- infectious A09.0
	- noninfectious K52.9
	- ulcerative (chronic) K51.1 K51.0
	Ileus (bowel) (colon) (inhibitory) (intestine) (neurogenic) K56.7 – meconium (with cystic fibrosis) E84.1+ P75*
Revise subterms:	- meaning meconium plug (without cystic fibrosis) P76.0
Revise subterms.	- without cystic fibrosis P76.0
	newborn
	- due to meconium (with cystic fibrosis) E84.1+ P75*
	meaning meconium plug (without cystic fibrosis) P76.0
	without cystic fibrosis P76.0
	– – transitory P76.1
Add lead term and	Immobility - due to prolonged bedrest R26.3
subterm; delete	- syndrome (paraplegic) M62.3
lead term	- Immobility syndrome (paraplegic) M62.3
Add lead term	Immobility syndrome (paraplegic) M62.3
	Immune compromised NEC D89.9
	Immunization (see also Vaccination) Z26.9
Add lead term	Immune compromised NEC D89.9 Immune reconstitution syndrome D89.3
	Immune compromised NEC D89.9
Add new lead	Immune compromised NEC D89.9 Immune reconstitution inflammatory syndrome [IRIS] D89.3
term	Immune reconstitution syndrome D89.3
Delete	Immunocytoma (M9671/3) C83.0
Delete	Ammunocy volum (1150 / 115) costo

INSTRUCTION	ALPHABETIC INDEX ENTRIES
morphology code	
	Impaired, impairment (function)
Revise code	- tolerance, glucose R73.0
Revise code	- vision NEC H54.7 <u>H54.9</u> - visual
Add subterms	binocular H54.9
	mild H54.3
	<u> moderate H54.2</u>
	<u> severe H54.1</u>
	<u> monocular</u> moderate H54.6
	severe H54.5
	Impaired, impairment (function)
Revise cross	- kidney (see also Failure, renal kidney) N19
reference and add	<u> acute N17</u>
subterms:	chronic N18.9
	<u> end-stage N18.5</u> stage 1 N18.1
	stage 2 N18.2
	stage 3 N18.3
	<u> stage 4 N18.4</u>
	<u> stage 5 N18.5</u>
	end-stage N18.5
	tubular function disorder N25.9 neonatal, transient P74.8
	- liver K72.9
Revise cross	
reference and add	- renal (see also Failure renal Impaired, impairment, kidney) N19
subterms:	— disorder resulting from N25.9
	- tolerance, glucose R73.0 Impaired, impairment (function)
Revise subterm	- kidney (see also Failure, renal-kidney) N19
Add modifier	Infarct, infarction (of)
	- intestine (acute) (agnogenic) (hemorrhagic) (nonocclusive) (transmural) K55.0
	- mesentery, mesenteric (embolic) (thrombotic) K55.0

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Add non-essential modifiers:	Infarct, infarction (of) - myocardium, myocardial (acute or with a stated duration of 4 weeks or less) I21.9 chronic or with a stated duration of over 4 weeks I25.8 healed or old I25.2 nontransmural I21.4 past (diagnosed on ECG or other special investigation, but currently presenting no symptoms) I25.2 - subsequent (extension) (recurrent) (reinfarction) I22.9 anterior (wall) I22.0
Revise codes	Infection, infected (opportunistic) B99
	alimentary canal NEC (<i>see also</i> Enteritis, infectious) A09 A09.0
	- colon (see also Enteritis, infectious) A09 A09.0
	- enteric (see also Enteritis, infectious) A09 A09.0
	- gastrointestinal (see also Enteritis, infectious) A09 A09.0
	- intestine, intestinal (<i>see also</i> Enteritis, infectious) A09 A09.0
Add subterms	Infection, infected (opportunistic) B99 - Burkholderiapseudomallei (see also Melioidosis) A24.4 NEC A49.8 - mallei A24.0
	as the cause of disease classified elsewhere B96.8 pseudomallei (see alsoMelioidosis) A24.4 as the cause of disease classified elsewhere B96.8
	- Pseudomonas, pseudomonad NEC A49.8
	as cause of disease classified elsewhere B96.5 mallei A24.0 - see Infection, Burkholderia, mallei
	pseudomallei (see also Melioidosis) A24.4 - <u>see Infection, Burkholderia, pseudomallei</u>
	Infection, infected (opportunistic) B99
	-bloodstream — see Septicaemia Sepsis
	- Clostridium, clostridium
Revise subterms:	difficile septicemia <u>sepsis</u> A41.4

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	gas-forming NEC A48.0
	perfringens as cause of disease classified elsewhere B96.7 foodborne (disease) A05.2 gas gangrene A48.0 septicaemia_sepsis_A41.4
	- generalised NEC (see also Septicaemia Sepsis) A41.9
	-pyemic — see Septicaemia Sepsis
	- Salmonella (arizonae) (cholerae-suis) (enteritidis) (typhimurium) A02.9 with
	(gastro)enteritis A02.0 septicemia sepsis A02.1
	specified manifestation NEC A02.8
	- septic
	generalised — see Septicaemia-Sepsis localised, skin (see also Abscess) L02.9
	localised, skill (<i>see diso</i> Abscess) Lo2.9 -septicemic — see Septicaemia-Sepsis
	-systemic — see Septicaemia Sepsis
	Infection, infected - continued
	- Cestodes – <i>see</i> Infestation, cestodes
Add subterm and	- <u>chest J22</u>
revise code:	- Chilomastix (intestinal) A07.8
	- lung (see also Pneumonia) 198.4 118.9
	Infection, infected (opportunistic) B99 Note: Parasitic diseases may be described as either 'infections' or 'infestations'; both lead terms should therefore be consulted.
	- Clostridium, clostridium
	- bifermentans A48.0
	- botulinum A05.1
	congenital P39.8
Revise code	– – difficile
	as cause of disease classified elsewhere B96.8

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	foodborne (disease) A05.8A04.7
	gas gangrene A48.0
	necrotizing enterocolitis A05.8-A04.7 septicemia A41.4
Add asterisk	Infection, infected (opportunistic) B99
codes	- Chlamydia, chlamydial A74.9
	− – genitourinary tract A56.2†N29.1*
	specified NEC A56.1 <u>†N29.1*</u>
Revise code	Infection, infected (opportunistic) B99
	- Helicobacter pylori, as cause of disease classified elsewhere <u>B96.8 B98.0</u>
Add subterms	Infection, infected (opportunistic) B99
	- Pseudomonas, pseudomanad NEC A49.8
	- pseudotuberculosis (extra–intestinal) A28.2
	– puerperal O86.4
Add subterms	Infection, infected, infective (opportunistic)
ridd Subterinis	– bleb, postprocedural H59.8
Add subterm	Infection, infected (opportunistic) B99
	_ vibrio vulnificus, as cause of disease classified elsewhere B98.1\
	Infection, infected (opportunistic) B99
	- with
	— antibiotic resistant bacterial agent (resistant to) U89.9
	Note: Use separate code to identify the infection. — methicillin U80.1
	— multiple antibiotics U88
Delete subterms	penicillin U80.0
	— penicillin related antibiotic U80.8
	— specified antibiotic (single) NEC U89.8
	—— multiple antibiotics U88
	— vancomycin U81.0
	- vancomycin related antibiotic U81.8
	Infection, infected (opportunistic) B99
Revise subterms	- Heterophyes (heterophyes) B66.8
	- hip (joint) NEC M00.9
	- due to internal joint prosthesis T84.5
	skin NEC L08.9

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	- joint see Arthritis, infectious NEC M00.9 - due to internal joint prosthesis T84.5 knee (skin)(joint)NEC-L08.9 M00.9 - due to internal joint prosthesis T84.5 — joint M00.9 - skin NEC L08.9 - shoulder (joint) NEC M00.9 - due to internal joint prosthesis T84.5 - skin NEC L08.9
Delete morphology code and cross reference and code	Infiltrate, infiltration – leukemic (M9800/3) see Leukemia – lymphatic (M9820/3) (see also Leukemia, lymphatic) C91.9 D47.9 – gland I88.9
Add subterms	Inflammation, inflamed, inflammatory (with exudation) - pleura – see Pleurisy - polyp - colon K51.4 - prostate (see also Prostatitis)
Revise code and add subterms	Inflammation, inflamed, inflammatory (with exudation) ileum (see also Enteritis) A09 A09.9 - infectious A09.0 - noninfectious K52.9
Add subterms	Inflammation, inflamed, inflammatory (with exudation) <u>bleb</u> , postprocedural H59.8
Revise subterms	Influenza (specific virus not identified) J11.1 - with - digestive manifestations J11.8 avian-certain identified influenza virus identified J09 other influenza virus identified J10.8 - enteritis J11.8 avian-certain identified influenza virus identified J09

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	other influenza virus identified J10.8
	gastroenteritis J11.8
	avian-certain identified influenza virus identified J09
	other influenza virus identified J10.8
	involvement of gastrointestinal tract J11.8
	gastromestman tract 311.6 = avian-certain identified influenza virus identified J09
	other influenza virus identified J10.8
	nervous system NEC J11.8
	avian certain identified influenza virus identified J09
	other influenza virus identified J10.8
	—— laryngitis J11.1
	avian -certain identified influenza virus identified J09
	other influenza virus identified J10.1
	manifestations NEC J11.8
	– – – avian-certain identified influenza virus identified J09
	other influenza virus identified J10.8
	– meningismus J11.8
	avian-certain identified influenza virus identified J09
	other influenza virus identified J10.8
	myocarditis J11.8† I41.1*
	avian certain identified influenza virus identified J09† I41.1*
	other influenza virus identified J10.8† I41.1*
	pharyngitis J11.1
	avian certain identified influenza virus identified J09 other influenza virus identified J10.1
	other influenza virus identified J10.1 pleural effusion NEC J11.1
	pieurai eriusion NEC 311.1 avian certain identified influenza virus identified J09
	atvian certain identified influenza virus identified J10.1
	- pneumonia (any form in J12-J16, J18) J11.0
	avian-certain identified influenza virus identified J09
	other influenza virus identified J10.0
	- – respiratory manifestations NEC J11.1
	avian certain identified influenza virus identified J09
	other influenza virus identified J10.1
	upper respiratory infection (acute) NEC J11.1
	avian certain identified influenza virus identified J09

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	 other influenza virus identified J10.1 - A/H1N1 [swine] J09
	- A/H5N1 [avian] J09
	- avian [A/H5N1] J09
	other J10.1
	– summer, of Italy A93.1
	<u>- swine [A/H1N1] J09</u>
	<u> other J10.1</u>
D 1.	- virus identified J10.1
Revise code	Inhibitor, systemic lupus erythematosus, presence D68.8-D68.6
	Injury (see also specified injury type) T14.9 - cord
Add non-essential	spermatic (pelvic region) S37.8
modifiers and	scrotal region S39.8
subterm:	- spermatic cord (scrotal pelvic region) \$39.9 \$37.8
	pelvic scrotal region \$\frac{\sample 337.8}{\sample 239.8}\$
Revise codes	Injury (see also specified injury type) T14.9
	- superficial (see also Contusion)
	– – finger S60.8 <u>S60.9</u>
	Injury (see also specified injury type) T14.9
	injury (see also specified injury type) 114.9
	- epiglottis NEC S19. 8 9
	- esophagus
	cervical NEC S19. 8 9
Revise codes	- gland
	thyroid NEC S19. <u>89</u>
	- larynx NEC S19.89
	- throat NEC S19.89 - thyroid (gland) NEC S19.89
	- trachea (cervical) NEC S19.89
	Injury (see also specified injury type) T14.9
Revise code	- periurethral tissue S37.3
	complicating delivery O71.5 <u>O70.0</u>
	Injury (see also specified injury type) T14.9

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Revise subterms and add cross	- epiglottis (see also Injury, neck)NEC S19.9
references	- esophagus (thoracic part) S27.8 cervical (<i>see also</i> Injury, neck) NEC S19.9
	- gland
	lacrimal (laceration) S05.8 salivary S09.9
	sanvary 509.9 thyroid (see also Injury, neck) NEC S19.9
	- larynx (see also Injury, neck)NEC S19.9
	mhowny (see also Injury neels)\$00,0\$10,0
	- pharynx (see also Injury, neck)\$09.9\S19.9
	- throat (see also Injury, neck)NEC S19.9
	thyroid (gland) (see also Injury, neck) NEC S19.9
	trachea (cervical) (see also Injury, neck)NEC S19.9
Add modifier and	Insufficiency, insufficient
subterms	- arterial I77.1
	mesenteric (chronic) K55.1
	<u> acute K55.0</u> - vascular I99
	- intestine K55.9
	mesenteric (chronic) K55.1
	<u> acute K55.0</u>
Revise subterms	Insufficiency, insufficient
	kidney (see also Failure, renal kidney) N19renal (see also Failure, renal kidney) N19
Revise code	Insulinoma (M8151/0)
	- pancreas D13.7
	<u> benign (M8152/0) D13.7</u>
	uncertain or unknown behavior (M8152/1) D37.7 malignant (M8152/3) C25.4
	malignant (M8152/3) C25.4 - specified site NEC—see Neoplasm, benign
	- benign (M8152/0) – see Neoplasm benign
	uncertain or unknown behavior (M8152/1) – see Neoplasm uncertain or unknown behavior

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	malignant (M8152/3) – see Neoplasm malignant
	- unspecified site D13.7
	benign (M8152/0) D13.7
	uncertain or unknown behavior (M8152/1) D37.7
	malignant (M8152/3) C25.4
	Intoxication - foodborne A05.9
	due to salmonella A02.9
	samonena A02.9
	(gastro)enteritis A02.0
	localised infection(s) A02.2
	septicemia sepsis A02.1
Revise subterms:	specified manifestation NEC A02.8
Revise subterms	Iridocyclitis H20.9
	– syphilitic (early) (secondary) A51.4† H22.0*
	congenital (early) A50.0† H22.0*
	late A50.3† H22.0*
	− − late A52.7† H22.0*
Revise subterms	Iritis (see also Iridocyclitis) H20.9
	– syphilitic (secondary) A51.4† H22.0*
	congenital (early) A50.0† H22.0*
	$\underline{ \text{late A}50.3\dagger \text{H}22.0*}$
	late A52.7† H22.0*
	Irritable, irritability R45.4
	heart (naveheanie) E45 2
Revise code:	- heart (psychogenic) F45.3 - hip M24.8 M65.8
Revise code.	- ileum K59.8
Add subterm	Ischemia, ischemic 199
rad subterm	
	- heart (chronic or with a stated duration of over 4 weeks) I25.9
	acute or with a stated duration of 4 weeks or less I24.9
	- <u>- failure I25.5</u>
Add subterm	Ischemia, ischemic 199
	- intestine (large) (small) K55.9
	acute K55.0

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	chronic K55.1
	- kidney N28.0
	- mesenteric, acute K55.0 chronic K55.1
Revise lead term	Jackson's membrane or veil-Q43.3
and add subterms	- membrane or veil Q43.3
and add subterms	– paralysis (syndrome) G83.8
Revise code	Jaundice (yellow) R17
	– fetus or newborn (physiological) P59.9
	due to or associated with
	hereditary hemolytic anemia <u>P58.8-D58.9</u>
Add non-essential modifiers and	Jaundice (yellow) R17
subterm	- familial nonhemolytic (congenital) (Gilbert) E80.4 - Criger-Najjar E80.5
Add code and	Jejunitis - see also Enteritis A09.9
subterm	- infectious A09.0
Delete non-	- noninfectious (presumed) K52.9
essential modifier	
	Jerks, myoclonic G25.3
Add lead term:	Jervell and Lange-Nielsen syndrome I45.8
	Jeune's disease Q77.2
Revise lead term	Kahler 's disease (M9732/3) C90.0
and delete	
morphology code Revise cod	Keratocyst (odontogenic) K09.0 D16.4
Revise code and	Keratocyst (odontogenic) (of) D16.4D16.5
add subterm	- upper jaw (bone) D16.4
Revise subterms	Keratoiritis (see also Iridocyclitis) H20.9
Revise subterins	- syphilitic (congenital late)(see also Iritis, syphilitc) A50.3† H22.0*
Add lead term	Kimmelstiel(-Wilson) disease or syndrome
and subterm	Kimura disease D21.9
	- specified site (see Neoplasm, connective tissue benign)
Revise code	Klatskin's tumour (M8162/3) C22.1 C24.0
Add subterms and	Labor (see also Delivery)
modify code	- early onset (before 37 completed weeks of gestation) O60
	<u> induced</u>

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	with delivery O60.3 without delivery O60.0 spontaneous with delivery preterm O60.1 term O60.2 without delivery O60.0 - premature or preterm induced with delivery O60.3 without delivery O60.0 - spontaneous with preterm delivery O60.1
	term delivery O60.2 without delivery O60.0
Revise codes and subterms	Laceration (see also Wound, open) T14.1 - periurethral tissue S37.8 - obstetric trauma O71.5-O70.0 - vagina (high) S31.4 - complicating delivery O71.4 - with perineum O70.0 - muscles O70.1 - vagina, vaginal wall (low) S31.4 - complicating delivery O70.0 and muscles (perineal)(vaginal) O70.1 high (deep)(instrumental)(mid)(sulcus) O71.4
Revise code:	Lambert-Eaton syndrome C80C80† G73.1*
Add cross reference	Lambert–Eaton syndrome (see also Neoplasm) C80† G73.1* – unassociated with neoplasm G70.8
Delete morphology code and revise code	Lennert's lymphoma (M9704/3) C84.34
Delete morphology code and revise code	Lesion (nontraumatic) – angiocentric immunoproliferative (M9766/1) D47.7 C83.8

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Revise code	Lesion (nontraumatic)
	••• ••• ••• ••• ••• ••• ••• ••• ••• ••
Delete code at	- angiocentric immunoproliferative <u>C83.8C86.0</u> Lesion (nontraumatic)
subterm level	
Subterin lever	- primary A51.0
	carate A67.0
	pinta A67.0
	syphilis A51.0
	yaws A66.0
Delete	Letterer–Siwe's disease (M9722/3) C96.0
morphology code Delete	Leukemia C95.9
morphology	- acute (bilineal) (biphenotypic) (mixed lineage) NEC (M9800/3) C95.0
codes, revise	- adult T-cell (M9827/3) C91.5
codes, and add	- aleukemic NEC (M9804/3) C95.79
subterms	<u>– B–cell type</u>
	<u>– lymphocytic, chronic C91.1</u>
	<u>– prolymphocytic, C91.3</u>
	<u>- basophilic (M9870/3) C92.7</u> <u>- acute C94.7</u>
	<u> acute C94.7</u> - bilineal, acute C95.0
	- biphenotypic, acute C95.0
	- blast (cell) (M9801/3) C95.0
	- blastic (M9801/3) C95.0
	– granulocytic (M9861/3) C92.9 <u>0</u>
	<u>- chronic NEC (M9803/3) C95.1</u>
	<u>- eosinophilic (M9880/3) C92.7</u>
	 - chronic [hypereosinophilic syndrome] D47.5 - erythroid, acute C94.0
	<u>erythroid, acute C94.0</u> – granulocytic (M9860/3) C92.9
	- acute (M9861/3) C92.0
	– aleukemic (M9864/3) C92. 7 9
	– blastic (M9861/3) C92.0
	– chronic (M9863/3) C92.1
	subacute (M9862/3) C92.2
	<u>hairy cell (M9940/3) C91.4</u>

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	 histiocytic (M9890/3) C93.9 lymphatic (M9820/3) C91.9
	<u>- lymphatic (M9820/3) C91.9</u> acute (M9821/3) C91.0
	- aleukemic (M9824/3) C91. 7 9
	chronic (M9823/3) C91.1
	subacute (M9822/3) C91.29
	- lymphoblastic (acute) [ALL] (M9821/3) C91.0
	<u>- lymphocytic (M9820/3) C91.9</u>
	acute (M9821/3) C91.0
	aleukemic (M9824/3) C91.79
	chronic (M9823/3) C91.1
	 - subacute (M9822/3) C91.29 - lymphoid NEC (M9820/3) C91.9
	- tymphoid NEC (M9821/3) C91.9 - acute (M9821/3) C91.0
	- aleukemic (M9824/3) C91.79
	$-$ blastic $\frac{(M9821/3)}{(M9821/3)}$ C91.0
	chronic (M9823/3) C91.1
	subacute (M9822/3) C91.29
	<u>– lymphoplasmacytic C91.1</u>
	– lymphosarcoma cell (M9850/3) C94.7 <u>C91.9</u>
	- mast cell (M9900/3) C94.3
	- megakaryoblastic (acute) C94.2
	- megakaryocytic (acute) (M9910/3) C94.2
	 mixed lineage, acute C95.0 monoblastic (acute)(monocytoid/monocytic) (M9891/3) C93.0
	- monocytic, monocytoid (M9890/3) C93.9
	- monocytic, monocytoid (M9891/3) C93.9 - acute (M9891/3) C93.0
	- aleukemic (M9894/3) C93.79
	- chronic (M9893/3) C93.1
	- Naegeli–type (M9863/3) C92.1 C93.1
	- subacute (M9892/3) C93.29
	— monomyelocytic, acute (M9867/3) — see Leukemia, myelomonocytic
	- myeloblastic (acute) (M9861/3) C92.0
	<u>1/ETO C92.0</u>
	<u>– M0 C92.0</u>
	<u>–– M1 C92.0</u>

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	$\frac{-M2 C92.0}{}$
	<u> M3 C92.4</u>
	<u> with t(15;17) and variants C92.4</u> <u> M4 C92.5</u>
	M4 Eo with inv(16) or t(16;16) C92.5
	$\frac{M4}{M5} \frac{10}{\text{C93.0}}$
	$\frac{1000000}{ \text{M5a C93.0}}$
	M5b C93.0
	- minimal differentiation C92.0
	<u>– NEC C92.0</u>
	<u>– (with maturation) C92.0</u>
	- with t(8;21) C92.0
	- myelocytic (M9860/3) C92.9
	acute (M9861/3) C92.0
	- chronic (M9863/3) C92.1
	 myelogenous (M9860/3) C92.9 acute (M9861/3) C92.0
	aleukemic (M9864/3) C92.79
	chronic (M9863/3) C92.1
	with crisis of blast cells C92.1
	Philadelphia chromosome (Phl) positive C92.1
	$\frac{1}{$
	- subacute (M9862/3) C92.2
	– myeloid (M9860/3) C92.9
	acute (M9861/3) C92.0
	with
	<u>11q23-abnormality C92.6</u>
	multilineage dysplasia C92.8
	<u> M6 (a) (b) C94.0</u> M7 C94.2
	<u> M7 C94.2</u> - aleukemic (M9864/3) C92.79
	chronic (BCR/ABL-positive [CML]) C92.1
	atypical, BCR/ABL-negative C92.2
	<u>– – NEC (without a FAB classification) C92.0</u>
	subacute (M9862/3) C92.2
	- myelomonocytic C92.5

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	acute (M9867/3) C92.5 chronic (CMML-1) (CMML-2) (with eosinophilia) (M9868/3) C92.7 C93.1 - juvenile C93.3 - Naegeli-type monocytic (M9863/3) C92.1 C93.1 - neutrophilic, chronic D47.1 - NK-cell, aggressive C94.7 - plasma cell (M9830/3) C90.1 - plasmacytic (M9830/3) C90.1 - plasmacytic (M9830/3) C90.1 - prolymphocytic (B-cell type) (M9825/3) C91.3 acute [PML] C92.4 - T-cell type C91.6 - promyelocytic, acute (M9866/3) C92.4 acute [PML] C92.4 - T-cell type C91.6 - stem cell (M9801/3) C95.0 - of unclear lineage C95.0
	 subacute NEC (M9802/3) C95.29 T-cell type adult (HTLV-1-associated) (acute) (chronic) (lymphomatoid) (smoldering) C91.5
	- large granular lymphocytic (associated with rheumatic arthritis) C91.7
	- prolymphocytic C91.6
	 thrombocytic (M9910/3) C94.2 undifferentiated (M9801/3) C95.0 unspecified cell type C95.9
	<u>– – acute C95.0</u>
	<u> chronic C95.1</u>
Delete morphology code and revise code	Leukosarcoma (M9850/3) C94.7 C85.9
Revise subterm, non-essential modifier and code	Lipiod – see also condition – histiocytosis (essential) D76.0 E75.2 — essential E75.2
Add lead term and	Lipochondrodystrophy E76.0 Lipodermatosclerosis I83.1

INSTRUCTION	ALPHABETIC INDEX ENTRIES
subterm:	- ulcerated I83.2
	Low
Add subterms	set ears Q17.4white blood cell count R72
Revise subterms	Low
Revise code	- function kidney (see also Failure, renal kidney) N19
Revise code	Lupus – continued - erythematosus (discoid)(local) L93.0
	systemic M32.9
	inhibitor (presence of) D68.8 D68.6
	Lymphadenitis
Revise code	- regional, nonbacterial A28.1 I88.8
Delete	Lymphadenopathy (generalized) R59.1
morphology code and revise code	– angioimmunoblastic (with dysproteinemia) (M9767/1) D47.7 C86.5
Delete	Lymphoblastoma (diffuse) (M9685/3) C83.5 C85.9
morphology code and revise code	
	Lymphocytic
	- chorioencephalitis(acute) (serous) A87.2† G05.1*
Add subterm	- choriomeningitis(acute) (serous) A87.2† G02.0* - colitis K52.8
Delete	Lymphogranuloma (malignant) (M9650/3) C81.9
morphology code	Y 1
Delete morphology code	Lymphogranulomatosis (malignant) (M9650/3) C81.9
Delete	Lymphoma (malignant) (M9590/3) C85.9
morphology	- adult T-cell (M9827/3) C91.5
codes, revise	- angiocentric T-cell (M9713/3) C85.7 C86.0
codes; add	- angioimmunoblastic [AILD] (M9705/3) C84.4 C86.5
subterms	- B-cell NEC (M9590/3) C85.1
	<u>– – diffuse large (ALK–positive) (anaplastic) (centroblastic) (immunoblastic) (plasmablastic) (T–cell rich) C83.3</u>
	<u>– extranodal, marginal zone of mucosa–associated lymphoid tissue (MALT–lymphoma) C88.4</u>
	<u>– – large</u>
	cell, mediastinal (thymic) C85.2

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	<u> intravascular C83.8</u>
	monocytoid (M9711/3) C85.7 <u>9</u>
	<u>– primary effusion C83.8</u>
	<u> </u>
	<u>B-CLL, non-leukemic variant C83.0</u>
	- B-precursor NEC (M9590/3) C85.1 C83.5
	- Burkitt's (atypical)(-like) (small noncleaved, diffuse) (undifferentiated) (M9687/3) C83.7
	- resulting from HIV disease B21.1
	 centroblastic (diffuse) (M9683/3) C83.83 follicular (M9697/3) C82.72
	- centroblastic-centrocytic (diffuse) (M9676/3) C83.89
	- centroblastic-centrocytic (diffuse) (M7676/3) C83.8 <u>2</u> - follicular (M9692/3) C82.75
	- centrocytic (M9674/3) C83. 8 1
	- cleaved cell (diffuse) (M9672/3) C83.1
	- with
	large cell, follicular (M9691/3) C82.1
	noncleaved, large cell (M9680/3) C83.39
	– follicular (M9695/3) C82. 0 9
	large (diffuse) (M9681/3) C83.3
	– – follicular (M9698/3) C82.2
	– small (diffuse) (M9672/3) C83.1
	- convoluted cell (M9685/3) C83.5
	– cutaneous <u>NEC (M9709/3)</u> C84.5 <u>8</u>
	– diffuse (M9595/3) C83.9
	- histiocytic (M9680/3) C83.3 <u>9</u>
	large cell (M9680/3) C83.3
	noncleaved (M9682/3) C83.3
	- lymphocytic (well differentiated) (M9670/3) C83.0
	intermediate differentiation (M9673/3) C83.80
	poorly differentiated (M9672/3) C83.1 C85.9
	- mixed cell type (M9675/3) C83.2 C85.9
	lymphocytic-histiocytic (M9675/3) C83.2 <u>C85.9</u>
	 small and large cell (M9675/3) C83.2 C85.9 - noncleaved (M9682/3) C83.39
	large cell (M9682/3) C83.3
	- - marge cell (M9686/3) C83.3 - - small cell (M9686/3) C83.0
L	– reticulum cell sarcoma (M9593/3) C83.3

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	small cell (M9670/3) C83.0 cleaved (M9672/3) C83.1 lymphocytic (M9670/3) C83.0 noncleaved, Burkitt's (M9687/3) C83.7 - follicle centre (centroblastic-centrocytic)
	<u> cutaneous C82.6</u> <u> diffuse C82.5</u>
	 follicular (centroblastic-centrocytic) (nodular) (with or without diffuse areas) (M9690/3) C82.9 centroblastic (M9697/3) C82.7 centrocytic (M9692/3) C82.7
	grade I C82.0 II C82.1 III NEC C82.2
	= - IIIa C82.3 = - IIIb C82.4 - histiocytic (M9698/3) C82.2
	 - large cell (cleaved) (noncleaved) (M9698/3) C82.2 - mixed cell type (M9691/3) C82.1 - noncleaved (large cell) (M9698/3) C82.23
	- small cleaved cell (M9695/3) C82.0 - and large cell (M9691/3) C82.1 - histiocytic (M9680/3) C83.3 C85.9
	 true (M9723/3) C96.38 Hodgkin's (M9650/3) C81.9 immunoblastic (diffuse) (large type) (M9684/3) C83.43 large cell (diffuse) (M9680/3) C83.3
	— with — small cell, mixed diffuse (M9675/3) C83.2 — small cleaved, mixed, follicular (M9691/3) C82.1
	anaplastic ALK-negative C84.7 ALK-positive C84.6
	CD30-positive C84.6 primary cutaneous C86.6

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	<u>– – diffuse C83.3</u>
	<u> </u>
	<u> </u>
	<u> small cleaved, mixed, follicular C82.1</u>
	<u> Ki-1+7 C84.6</u>
	<u>– – noncleaved and cleaved C83.3</u>
	— Ki 1+ (M9714/3) C85.7
	— noncleaved and cleaved (M9680/3) C83.3
	- Lennert's (M9704/3) C84.34
	 leukemia, adult T-cell (M9827/3) C91.5 lymphoblastic (diffuse) (M9685/3) C83.5
	- lymphocytic (diffuse) (M9670/3) C83.0
	- intermediate differentiation, nodular (M9694/3) C82.79
	- nodular (M9690/3) (intermediate differentiation)(poorly differentiated)(well differentiated) C82.9
	intermediate differentiation (M9694/3) C82.7
	poorly differentiated (M9696/3) C82.7
	— — well differentiated (M9693/3) C82.7 — — small (M9670/3) C83.0
	- Sman (M97070/3) C83.0 - lymphoepithelioid (M9704/3) C84. 3 4
	- lymphoid tissue
	- bronchial-associated (BALT-lymphoma) C88.4
	<u>– – skin–associated (SALT– lymphoma) C88.4</u>
	<u>- lymphoplasmacytic C83.0</u>
	with, IgM-production C88.0
	 lymphoplasmacytoid (M9671/3) C83.80 lymphoplasmatic (M9671/3) C83.80
	- mantle zone (M9673/3) C83.8
	cell <u>C83.1</u>
	<u>– – zone C83.1</u>
	– Mediterranean (M9764/3) C88.3
	- mixed cell type
	 – diffuse (M9675/3) C83.2 C85.9 – follicular (M9691/3) C82.1
	- Ioniculai (M9091/3) Co2.1 - Iymphocytic–histiocytic (diffuse) (M9675/3) C83.2 <u>C85.9</u>
	nodular (M9691/3) C82.1

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	 - small and large cell (diffuse) (M9675/3) C83.2 C85.9
	- small cleaved and large cell, follicular (M9691/3) C82.1
	- monocytoid B-cell (M9711/3) C85. 7 9
	- NK-cell, blastic C86.4
	- nodal marginal C83.0
	- nodular (with or without diffuse areas) (M9690/3) C82.9
	 – histiocytic (M9698/3) C82.2 – lymphocytic (M9690/3) C82.9
	- - lymphocytic (W9694/3) C82.9 - - intermediate differentiation (M9694/3) C82.79
	poorly differentiated (M9696/3) C82.79
	$-$ - well differentiated $\frac{\text{(M9693/3)}}{\text{C82.79}}$ C82.79
	$-$ mixed (cell type) $\frac{(M9691/3)}{C82.12}$
	- mixed lymphocytic–histiocytic (M9691/3) C82.1
	- non-Burkitt 's , undifferentiated cell (M9686/3) C83.63
	- noncleaved (diffuse) (M9682/3) C83.39
	follicular (M9698/3) C82.2
	large cell (diffuse) (M9682/3) C83.3
	– – follicular (M9698/3) C82.2
	 - small cell (diffuse) (M9686/3) C83.0
	- non-Hodgkin's- (type) NEC-(M9591/3)- C85.9
	<u>– non-follicular (diffuse) NEC C83.9</u>
	 resulting from HIV disease B21.2
	<u>– (type) NEC C85.9</u>
	– peripheral T–cell (M9702/3) C84.4
	AILD (M9705/3) C84.4 <u>C86.5</u>
	angioimmunoblastic lymphadenopathy with dysproteinemia (M9705/3) C84.4 C86.5
	pleomorphic
	medium and large cell (M9707/3) C84.4
	small cell (M9706/3) C84.4
	- plasmacytic (M9671/3) C83.80
	- plasmacytic-lymphocytic C83.0
	 plasmacytoid (M9671/3) C83.80 small lymphocytic C83.0
	- resulting from HIV disease B21.2
	- resulting from FTV disease B21.2 - small cell (diffuse) (M9670/3) C83.0
	- smart ceri (dirituse) (M7676/3) Co.3.0 - with large cell, mixed (diffuse) (M9675/3) C83.2 C85.9
	- cleaved (diffuse) (M9672/3) C83.1
	ciourea (unitato) (11701213) COST

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	and large cell, mixed, follicular (M9691/3) C82.1 follicular (M9695/3) C82.09 - lymphocytic (diffuse) (M9686/3) C83.0 - noncleaved (diffuse) (M9686/3) C83.0 - noncleaved (diffuse) (M9686/3) C83.7 - T - Cell NEC (M9590/3) C84.54 - adult (M9827/3) C91.5 - angioentric (M9713/3) C85.7 C86.0 - angioimmunoblastic C86.5 - cutaneous C84.8 - enteropathy associated C86.2 - enteropathy-type (intestinal) C86.2 - hepatosplenic (alpha-beta and gamma-delta types) C86.1 - mature, not classified C84.4 - panniculitis-like, subcutaneous C86.3 - peripheral AILD(M9702/3) C84.45 C86.5 - AILD (M9705/3) C84.4 - angioimmunoblastic lymphadenopathy with dysproteinemia (M9705/3) C84.4 C86.5 pleomorphic medium and large cell (M9707/3) C84.4 - primary cutaneous, CB30-positive C86.6 - T-precursor NEC C83.5 - T/NK-cell NEC C84.5 - extranodal, nasal type C86.0 - mature NEC C84.9 - true histiocytic (M9723/3) C96.38
	 T-zone (M9703/3) C84.24 undifferentiated cell (M9686/3) C83.69 Burkitt's type (M9687/3) C83.7
Delete morphology code	Lymphomatosis (M9590/3) – see Lymphoma
Delete morphology codes and revise codes	Lymphosarcoma (diffuse) (M9592/3) C85.09 - cell leukemia (M9850/3) C94.7 C91.9 - follicular (M9690/3) C82.9 - mixed cell type (M9691/3) C82.49

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Revise subterm and delete morphology code	Macroglobulinemia (idiopathic) (primary) C88.0 – Waldenström's (M9761/3) C88.0
Add subterm	Macroglobulinemia (idiopathic) (primary) C88.0 - monoclonal (essential) D47.2 - Waldenström C88.0
Revise code:	Malposition - congenital - aorta Q25.4 - arterial trunk Q25.4 Q20.0
Revise code	Malposition congenital
Add subterm	eustachian tube Q17.8 Q16.4 Management (of) - cardiac pacemaker NEC Z45.0 - implanted device Z45.9 - cardiac Z45.0 - specified NEC Z45.8
Revise code	March - fracture \$92.3 M84.3
Revise code	Marrow (bone) - arrest D61.9 - poor function D75.8 D61.9
Delete morphology code and revise code	Mast cell - disease, systemic tissue (M9741/3) C96.2 D47.0 - leukemia (M9900/3) C94.3 - sarcoma (M9740/3) C96.2 - tumor (M9740/1) D47.0 - malignant (M9740/3) C96.2
Delete morphology code	Mastocytoma (M9740/1) D47.0 – malignant (M9740/3) C96.2
Delete morphology code and add subterms	Mastocytosis Q82.2 - malignant (M9741/3) C96.2 - aggressive C96.2

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	<u>- systemic</u> (associated with clonal haematopoietic non-mast-cell disease) (indolent) (SM-AHNMD) D47.0
Add non-essential modifiers and add subterms:	Meconium - ileus (with cystic fibrosis), fetus or newborn E84.1+ P75* - meaning meconium plug P76.0 - without cystic fibrosis P76.0 - obstruction, fetus or newborn P76.0 - in mucoviscidosis E84.1+ P75* - plug syndrome (newborn) NEC P76.0
Add subterm and	Megacolon (acquired) (functional) (not Hirschsprung's disease) (in) K59.3
modify subterm	- toxic <u>NEC</u> K59.3 due to Clostridium difficile A04.7
Revise code:	Melanoma—continued - metastatic - specified site NEC (M8720/6) C79.8 - unspecified site (M8720/6) C80C79.9
Revise code.	— unspectified site (W6720/0)
Revise subterms:	pulmonary (chronic) A24.2 acute A24.1 subacute A24.2 - septicemia sepsis A24.1
Revise subterms	Meningismus R29.1 - due to serum or vaccine R29.1 - influenzal (specific virus not identified) J11.8 avian_certain identified influenza virus identified J09 - other influenza virus identified J10.8
	Meningitis (basal) (cerebral) (spinal) G03.9
Revise code:	- sterile G03.90 Meningoencephalitis (see also Encephalitis) G04.9
Revise codes	– pneumococcal G00.1 G04.2
Add lead term and	Mesenchymoma (M8990/1) – see also
subterm	Mesentery, mesenteric - see condition Mesenteritis K65.9

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	- sclerosing K65.8
Modify subterm and add code	Metaplasia - apocrine (breast) N60.8 R87 - cervix (squamous) — see Dysplasia, cervix R87
Revise code	Metaplasia - apocrine (breast) R87 N60.8
	Metastasis, metastatic
Revise code:	– cancer or neoplasm (M8000/6) C80 <u>C79.9</u>
Revise non-	Metatarsus, metatarsal (see also condition)
essential	- valgus (adductus) (abductus), congenital Q66.6
modifiers	- varus (adductus) (congenital) Q66.2 Microglioma (M9594/3) C85.7
Delete morphology code	
Add index lead	Millar's asthma J38.5
term:	Miller Fisher Syndrome G61.0
Add lead term and	Mucositis (drug induced) (radiation induced) (ulcerative) – (see also
subterms	Inflammation, by site) K12.3 – anus, anal canal K92.8
	- colon K92.8
	- duodenum K92.8
	- esophagus K92.8
	- gastric K92.8
	<u>– gastrointestinal (ulcerative) K92.8</u>
	<u>– ileum K92.8</u>
	- intestine (any part) K92.8
	- mouth K12.3 - nasal (ulcerative) J34.8
	- nasar (urcerative) 334.8 - oral K12.3
	- oropharyngeal K12.3
	– perianal K92.8
	<u>– perirectal K92.8</u>
	<u>- rectum K92.8</u>
	<u>- sigmoid K92.8</u>
	- stomach K92.8
	– vagina N76.8

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	<u>– vulva N76.8</u>
Add non-essential modifier	Mucinosis (cutaneous) (focal) (papular) (reticular erythematous) (skin) L98.5
Add lead term and subterm	Mutation - prothrombin gene [factor V Leiden mutation] D68.5
	Myasthenia, myasthenic G70.9 — malignant neoplasm NEC (M8000/3) (see also Neoplasm, malignant) C80C80† G73.2*
Revise code: Delete morphology code	Mycosis, mycotic NEC B49 – fungoides (M9700/3) C84.0
Delete morphology code and add non– essential modifiers	Myelodysplasia (M9989/1) (related to alkylating agent)(related to Epipodophyllotoxin)(related to therapy) – specified NEC D46.7 – spinal cord (congenital) Q06.1
Delete morphology code, add non–essential modifiers, revise code and add subterm	Myelofibrosis (chronic)(idiopathic)(with myeloid metaplasia) (M9961/1) D47.14 - acute (M9932/3) C94.54 - secondary, in myeloproliferative disease D47.4
Delete morphology code and revise code	Myeloma (multiple) (M9732/3) C90.0 - monostotic (M9731/3) C90.23 - plasma cell (M9732/3) C90.03 - plasma cell C90.3 - solitary (M9731/3) C90.23
Revise codes at subterms	Myeloma (multiple) C90.0 - monostotic C90.3 - plasma cell C90.3 C90.0 - plasma cell C90.3 C90.0
Delete morphology code	Myelomata, multiple (M9732/3) C90.0
Delete morphology code	Myelomatosis (M9732/3) C90.0
Delete	Myelosarcoma (M9930/3) C92.3

INSTRUCTION	ALPHABETIC INDEX ENTRIES						
morphology code							
Delete morphology code, revise codes	Myelosclerosis D75.8 D47.4 - with myeloid metaplasia (M9961/1) D47.14 - disseminated, of nervous system G35 - megakaryocytic (with myeloid metaplasia) (M9961/1) D47.14						
Delete morphology codes and revise codes	Myelosis - acute (M9861/3) C92.0 - aleukemic (M9864/3) C92.79 - chronic (M9863/3) C92.1 D47.1 - erythremic (M9840/3) C94.0 - acute (M9841/3) C94.0 - megakaryocytic (M9910/3) C94.2 - nonleukemic D72.8 - subacute (M9862/3) C92.29						
Revise subterms	Myocarditis (chronic) (fibroid) (interstitial) (old) (progressive) (senile) (with arteriosclerosis) I51.4 – influenzal (specific virus not identified) J11.8† I41.1* – avian certain identified influenza virus identified J09† I41.1* – other influenza virus identified J10.8† I41.1*						
Revise code	Myoma (M8895/0) — see also Neoplasm, connective tissue, benign - malignant (M8895/3) — see Neoplasm, connective tissue, malignant - prostate N40-D29.1 - uterus (cervix) (corpus) — see Leiomyoma						
Pavisa andar	Myopathy—continued in (due to) malignant recorder NEC (M8000/2) (see also Neopleam malignant) C80C80 # M62.8*						
Revise code: Add subterms	 – malignant neoplasm NEC (M8000/3) (see also Neoplasm, malignant) C80C80† M63.8* Myositis 						
7 Aud Subtellis	in (due to) trichinellosis B75† M63.1 * - inclusion body (IBM) G72.4 - infective M60.0 - interstitial M60.1 - juvenile M33.0						

INSTRUCTION	ALPHABETIC INDEX ENTRIES						
	- multiple – see Polymyositis						
Delete	Naegeli's						
morphology code	- disease Q82.8						
and revise code	– leukemia, monocytic (M9863/3) C92.1 <u>C93.1</u>						
Revise	Necrosis, necrotic (ischaemic) (see also Gangrene) R02						
Add subterms	- pressure L89 <u>.2</u>						
Add subternis	stage						
	<u> III L89.2</u> IV L89.3						
Add subterms and	Neglect (newborn) T74.0						
codes	- emotional, in childhood Z62.4						
	<u>- self R46.8</u>						
	causing insufficient intake of food and water R63.6						
Add subterm	Neonatal – see also condition						
	- abstinence syndrome P96.1 - tooth, teeth, K00.6						
	Neoplasm, neoplastic						
	- bone						
Revise code and	vomer C40.0 <u>C41.0</u> C79.5 - D16.4 D48.0						
add subterm:							
	<u>- vomer C41.0 C79.5 - D16.4 D48.0</u>						
	Malignant Jncertain or						
	Primary Secondar In situ Benign unknown behavior						
	y						
	Neoplasm, C80C80.9 C80C79.9 D09.9 D36.9 D48.9 neoplastic						
	p. 412						
	– disease, generalized C80						
	primary site not indicated C80.9 C79.9						
	primary site unknown, so stated C80.0 C79.9						
Add subterms and	Neoplasm, neoplastic—continued						
codes:	- disseminated C80						
	- primary site not indicated C80.9 C79.9						
	<u>– primary site unknown, so stated</u> <u>C80.0</u> <u>C79.9</u>						

INSTRUCTION	ALPHABETIC INDEX ENTRIES							
	 generalized primary site not inc primary site unkno metastatic, primary site primary site not inc 	wn, so stat		C80.9 C80.9 C80.0	<u>C79.9</u> <u>C79.9</u> <u>C79.9</u>			
	 – primary site unknown primary site unknown unknown site so state 	wn, so state	e <u>d</u>	C80.0 C80.0 C80.0	C79.9 C79.9 C79.9			
Revise code:	- gallbladder	C22.1 C7 C23 C7	8. 8 7 D01.5 D 8.8 D01.5 D1	13.4 D37.6	Ď			
Revise cross reference	Neoplasm, neoplastic – – central – nervous system – see			em				
Revise codes	Neoplasm, neoplastic – – parasellar			4 <u>3</u> D	33. <u>92</u> D4:	3. <u>92</u>		
Revise codes	Neoplasm, neoplastic—continued - septum— continued - rectovaginal							
	Neoplasm, neoplastic	•••						
	Neoplasm, neoplastic		Malignant secondary	Insitu	Benign	Unknown Uncertain behaviour		
Delete code	- lymph, lymphatic							
	gland (secondary)	-	C77.9	_	D36.0	D48.7		
	Virchow's	-	C77.0	-	D36.0	D48.7		
	- Virchow's gland Neoplasm, neoplastic	C77.0	C77.0	-	D36.0	D48.7		
Revise Code	- sphincter							

INSTRUCTION	ALPHABETIC INDEX ENTRIES								
	of Oddi								
		Malignant primary	Malignant secondary	Insitu	Benign	Unknown Uncertain behaviour			
	- bile or biliary (tract)	C24.9	C78.8	D01.5	5 D13.5	D37.6			
	canals, interlobular	C22.1	C78. 8- 7	D01.5	5 D13.4	D37.6			
Revise codes	duct or passage (common) (cystic) (extrahepatic)	C24.0	C78.8	D01.5	5 D13.5	D37.6			
	interlobular	C22.1	C78. 8 -7	D01.5	5 D13.4	D37.6			
	intrahepatic	C22.1	C78. 8- 7	D01.5	5 D13.4	D37.6			
	- canaliculi								
	biliferi	C22.1	C78. 8- 7	D01.5	D13.4	D37.6			
	intrahepatic	C22.1	C78. 8- 7	D01.5	5 D13.4	D37.6			
	- cholangiole	C22.1	C78. 8 - <u>7</u>	D01.5	5 D13.4	D37.6			
		t	Iı	ı situ					
		Primary	Primary Secondary						
	Neoplasm, neoplastic	C80.9	C79.9	D	009.9				
	- disease, generalized		C79.9						
	primary site not indicated	C80.9	C79.9						
Delete subterms:	primary site unknown, so stated	C80.9	C79.9						
	- disseminated		<u>C79.9</u>						
	- primary site not indicated	C80.9	C79.9						
	primary site unknown, so stated	C80.9	C79.9						
	- generalized		<u>C79.9</u>						
	primary site not indicated		C79.9						
	primary site unknown, so	C80.9	C79.9						

INSTRUCTION	ALPHABETIC INDEX ENTRIES						
	stated						
	- metastatic (multiple)		<u>C79.9</u>				
	— primary site not indicated	C80.9	C79.9				
	primary site unknown, so stated	C80.9	C79.9				
	- multiple, malignant (independent primary sites) <u>NEC</u>	C97 <u>C80.9</u>					
	independent primary sites	<u>C97</u>					
	secondary NEC		<u>C79.9</u>				
	- primary site unknown, so stated	C80.0	C79.9				
	- unknown site, primary	C80					
	- unknown site, whether prim	ary or second	lary C80.9				
Add row and revise code in	Neoplasm, neoplastic						
Neoplasm table	- urethrovaginal (septum)	C57.9 C	79.8 D07.3	D28.9 I	D39.9		
r (Copingin thoi	septum		279.8 D07.3	D28.9 I	D39.7		
	- vesicovaginal		79.8 D07.3		D39.9		
	septum	C57.9 C	279.8 D07.3	D28.9 I	D39. 9 7		
Add subterm	Neoplasm						
	•••	~~. ~ ~					
Revise codes	- hepatobiliary	C24.9 C	78.8 D01.5	D13.9 I	<u> </u>		
Revise codes	Neoplasm, neoplastic - eyeball	C69.49 C	C79.4 D09.2	D31.49 I	D48 7		
	- intraocular	C69.49 C					
Revise codes	Neoplasm, neoplastic - coccygeal	_					
		75.5 <u>C49.5</u> (C79.8 D35.6 <u>D</u>	21.5 D44.	7 <u>D48.1</u>		
	- glomus	275 5040 5	C79.8 - D35.6	D21.5 D44	7D49 1		
	coccygeal • Nephritis, nephritic N05	. 13.3 <u>C49.3</u>	C17.0 - D33.0	<u>D21.3</u> D44.	7 <u>U40.1</u>		
	• •						
	••••						

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Revise cross reference:	- with - glomerular lesion - diffuse sclerosing (see also Failure, renal, Disease, kidney, chronic) N18.9 - hypocomplementemic — see Nephritis, membranoproliferative
	- degenerative — see Nephrosis - diffuse sclerosing (see also Failure, renal, Disease, kidney, chronic) N18.9
	- sclerosing, diffuse (see also Failure, renal Disease, kidney, chronic) N18.9
Add subterm	Nephritis, nephritic N05
	- cirrhotic (see also Sclerosis, renal) N26 - complicating pregnancy, childbirth or the puerperium O99.8-O26.8
Add subterm	Nephrolithiasis (congenital)(pelvis)(recurrent) (see also Calculus, kidney) N20.0
	- congenital Q63.8
	- gouty M10.0† N22.8*
	Nephropathy (see also Nephritis) N28.9
	Note: Where a term is indexed only at the three character level, eg N07, reference should be made to the list of fourth character subdivisions in Volume 1 at N00-N08.
	- chemical — see Nephropathy, toxic
	- diabetic (see also E10-E14 with fourth character .2) E14.2† N08.3*
	- drug-induced N14.2
	specified NEC N14.1 - focal and segmental hyalinosis or sclerosis N02.1
	- heavy metal-induced N14.3
Add subterms:	- hereditary NEC N07
Tide succession.	- hypertensive (see also Hypertension, kidney) I12.9
	end-stage (failure) I12.0 - IgA N02.8
	- with glomerular lesion N02
	specified pathology NEC N02.8
	- lead N14.3 - membranoproliferative (diffuse) N02.5
	memoranopromerative (unituse) 1302.5

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	- membranous (diffuse) N02.2
	- mesangial (IgA/IgG) — see Nephropathy, IgA
	proliferative (diffuse) N02.3 - mesangiocapillary (diffuse) N02.5
	- obstructive N13.8
	Neuritis M79.2
	- progressive hypertrophic interstitial G60.0
Revise code:	– puerperal, postpartum O26.8 <u>O90.8</u>
D 1	- retrobulbar H46
Revise subterm	Neuritis – arising during pregnancy O26.8
	– peripheral (nerve) G62.9
	– complicating pregnancy or puerperium O26.8
	– pregnancy-related O26.8
	Neuropathy, neuropathic—continued
Revise code:	– carcinomatous C80 C80† G13.0*
Revise code . Revise subterms	Neuropathy, neuropathic
Revise subterms	- uremic N188-5 † G63.8*
Add non-essential	Nodule(s), nodular
modifier and	- thyroid (gland) E04.1
subterm	toxic or with hyperthyroidism (multinodular) E05.2
	<u> single E05.1</u>
Revise cross	Non-functioning – kidney (see also Failure, renal kidney) N19
reference	Non-Hodgkin's lymphoma NEC (M9591/3) C85.9
Delete morphology code	Non-Hougkin-s lymphoma NEC (1819391/3) C63.9
and revise lead	
term	
	Obstruction, obstructed, obstructive
	- intestine (mechanical)(neurogenic)(paroxysmal)(postinfective)(reflex) K56.6
	congenital (small) Q41.9
	large Q42.9 specified part NEC Q42.8
	specified part NEC Q42.0

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Add subterm	<u>– – distal (syndrome) E84.1</u>
Revise code at subterm	Obstruction, obstructed, obstructive - airway J98.8 chronic J44.9 with allergic alveolitis J67.9 asthma NEC J45.9 J44
Add subterm	Obstruction hepatobiliary K83.1
Add modifier and	Occlusion, occluded
subterm	- artery — see also Embolism, artery - mesenteric (chronic) K55.1 acute K55.0 - mesenteric artery (chronic) K55.1 - acute K55.0
Delete code at	Orthodontics Z51.8
lead term	adjustment Z46.4fitting Z46.4
Revise code	Osteomyelofibrosis D75.8 <u>D47.4</u>
Revise code	Osteomyelosclerosis <u>D75.8 D47.4</u>
Add subterm	Osteonecrosis _ jaw (drug-induced)(radiation-induced) K10.2
Delete	Osteoporosis M81.9
morphology code	- in (due to) - multiple myelomatosis (M9732/3) C90.0† M82.0*
Revise code	Osteosclerosis (fragilitas)(generalisata) Q78.2 – myelofibrosis D75.8 D47.4
Revise subterms	Otitis—continued - media—continued - in (due to) influenza (specific virus not identified) J11.8† H67.1* avian certain identified influenza virus identified J09† H67.1* other virus identified J10.8† H67.1*

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	Pain(s) R52.9
Revise code	- shoulder <u>M75.8M25.5</u>
Revise code and	Palsy (see also Paralysis) G83.9
subterm	- bulbar (chronic) (progressive) G12.2
	- of childhood (Fazio-Londe) G12.2
	- pseudo NEC G12.2
	— supranuclear NEC G12.2
	- supranuclear (progressive) G23.1
Add lead term	Pancolitis, ulcerative (chronic) K51.0
Delete	Panmyelosis (acute) (M9931/3) C94.4
morphology and	<u>with myelofibrosis, acute C94.4</u>
add subterm	Panniculitis M79.3
Add subterms	- febrile nodular nonsuppurative (Weber-Christian) M35.6
	- nodular, nonsuppurative W79.3 M35.6
	<u>relapsing</u> (febrile nodular nonsuppurative)(Weber-Christian) M35.6
Add lead term	Papulosis, lymphomatoid C86.6
Revise code	Papulosis, lymphomatoid <u>L41.2C86.6</u>
Revise asterisk	Paracoccidioidomycosis B41.9
code:	– pulmonary B41.0 † J99.8 <u>J17.2*</u>
Revise lead term,	Paragranuloma, Hodgkin's (M9660/3) C81.7 – Hodgkin C81.0
delete morphology code	- nodular C81.0
and add subterms	
Revise code	Parakeratosis R23.4
Revise code	– variegata <u>L41.0</u> <u>L41.5</u>
Modify	Paralysis, paralytic (complete) (incomplete) G83.9
	- agitans (see also Parkinsonism) G20
Revise code	arteriosclerotic G21.8 G21.4
	Paralysis, paralytic (complete) (see also Paresis) G83.9
Revise code:	 - uremic N18. 8 5† G99.8*
	Paralysis, paralytic (complete) (incomplete) (see also Paresis) G83.9
Add dagger	

INSTRUCTION	ALPHABETIC INDEX ENTRIES
symbol, asterisk	- throat J39.2
code and revise	diphtheritic A36.0 <u>† G99.8*</u>
code	– Weber's I67.9 <u>I67.8†</u> G46.3*
Revise code and	Paralysis, paralytic- continued
subterm	
	- bulbar (chronic) (progressive) G12.2
	infantile (see also Poliomyelitis, paralytic) A80.3
	poliomyelitic (<i>see also</i> Poliomyelitis, paralytic) A80.3
	pseudo NEC G12.2 — supranuclear NEC G12.2
	progressive (atrophic) (bulbar) (spinal) G12.2
	supranuclear G23.1
	- supranuclear (progressive)G12.2 G23.1
Delete	Paraproteinemia D89.2
morphology codes	– benign (familial) D89.2
	- monoclonal (M9765/1) D47.2
	– secondary to malignant disease (M9765/1) D47.2
	Parkinsonism (idiopathic) (primary) G20
A 11 1.	- with orthostatic hypotension (idiopathic) (symptomatic) G90.3
Add subterm	- arteriosclerotic G21.4
	- secondary G21.9 due to
Revise code	due to arteriosclerosis G21.8 <u>G21.4</u>
Revise code	drugs NEC G21.1
	neuroleptic G21.0
	encephalitis G21.3
	external agents NEC G21.2
	syphilis A52.1† G22*
	specified NEC G21.8
	- syphilitic A52.1† G22*
	- treatment-induced NEC G21.1
Add subterm	- vascular G21.4
	Parkinsonism (idiopathic) (primary) G20

ALPHABETIC INDEX ENTRIES								
- with <u>neurogenic</u> orthostatic hypotension (idiopathic) (symptomatic) G90.3								
Parulis K04.67								
- with sinus K04.6								
Perforation, perforated (nontraumatic) - accidental during procedure (blood vessel) (nerve) (organ) T81.2 - appendix K35.0 with peritonitis, generalized K35.2 peritonitis, localized K35.3 - bile duct (common) (hepatic) K83.2								
Pericarditis (with decompensation) (with effusion) I31.9 - uremic N18.85† I32.8*								
Perihepatitis K65.8 -chlamydial A74.8† K67.0* - gonococcal A54.8† K67.1*								
Peritonitis (adhesive) (fibrinous) (with effusion) K65.9 - with or following - abscess K65.0 - appendicitis K35.9 generalized K35.2 with mention of perforation or rupture K35.0 K35.2 - localized K35.3 with mention of perforation or rupture K35.3 - diverticular disease (intestine) K57.8								
Peritonitis (adhesive) (fibrinous) (with effusion) K65.9 - due to foreign substance accidentally left during a procedure (chemical)(powder) (talc) T81.6 - eosinophilic K65.8 acute K65.0								
Peritonitis(adhesive) (fibrinous) (with effusion) K65.9 - with or following - appendicitis K35.9K35.3 — generalized K35.2 — with mention of perforation or rupture K35.2 — localized K35.3 — with mention of perforation or rupture K35.3								

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	- diffuse (see also Peritonitis, generalized) K65.0 - disseminated (see also Peritonitis, generalized) K65.0 - general(ized) K65.0 - with or following appendicitis (acute) (perforation) (rupture) K35.2 - distributed (see also Peritonitis, generalized) K65.0 - with or following appendicitis (acute) (perforation) (rupture) K35.2 - distributed (see also Peritonitis, generalized) K65.0
	- localized K65.0 - with or following appendicitis (acute) (perforation) (rupture) K35.3
Revise code and add subterm	Phlegmasia - alba dolens (puerperal) O87.1 - complicating pregnancy O22.3 - nonpuerperal nonobstetric I80.1 - cerulea dolens I80.2
	Placenta
Revise code Revise subterms	- syphilitic A52.7-098.1 Placenta, placental
and codes	- accreta (with hemorrhage) O72.0 O43.2 - without hemorrhage O73.0 - adherent (with postpartum hemorrhage) (morbidly) O72.0 O43.2 - without hemorrhage O73.0 - increta (with postpartum hemorrhage) O72.0 O43.2 - without hemorrhage O73.0 - membranacea O43.1 - morbidly adherent O43.2 - multilobed O43.1 - necrosis O43.8 - percreta (with postpartum hemorrhage) O72.0 O43.2 - without hemorrhage O73.0
Delete morphology code and non–essential modifiers, add subterms, revise code	Plasmacytoma (extramedullary)(solitary) (M9731/3) NEC C90.23 - extramedullary C90.2 - medullary C90.0 - solitary C90.3

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Revise	Plaster ulcer L89
Add subterms	- stage
	<u>IL89.0</u>
	<u> II L89.1</u>
	<u> III L89.2</u>
	- <u>IV L89.3</u>
Revise code	Pleurisy (acute) (adhesive) (chronic) (double) (dry) (fibrinous) (subacute) R09.1
	- staphylococcal J86.9 <u>J90</u>
Modify	- streptococcal J90
Widdify	Pneumonia (acute) (double) (migratory) (purulent) (septic) (unresolved) J18.9 - interstitial J84.9
Add subterm	- interstitian 384.9 usual J84.1
	Pneumonia (acute) (double) (migratory) (purulent) (septic) (unresolved) J18.9
	- in (due to)
	schistosomiasis B65† J17.3*
Revise subterms:	septicemia sepsis A41† J17.0*
Add subterm	Pneumonia
	- cholesterol J84.8
	- chronic J98.4
A 11 1.	- cirrhotic (chronic)(see also Fibrosis, lung) J84.1 Pneumonia (acute) (double) (migratory) (purulent) (septic) (unresolved) J18.9
Add subterms	- in (due to)
	- histoplasmosis B39.2 †J17.2*
	– human metapneumovirus J12.3
	Pneumonia (acute) (double) (migratory) (purulent) (septic) (unresolved) J18.9
Add non–essential	
modifier and	– alba (early congenital syphilitic) A50.0† (J17.0*)
asterisk code	
	– syphilitic, congenital early (early) A50.0† (J17.0*)
	Pneumonia (acute) (double) (migratory) (purulent) (septic) (unresolved) J18.9
	- with
Revise subterms	- influenza, flu, or grippe (specific virus not identified) J11.0
	– – avian certain identified influenza virus identified J09
	other influenza virus identified J10.0

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	Polioencephalitis
Revise subterms	 influenzal (specific virus not identified) J11.8† G05.1* avian certain identified influenza virus identified J09† G05.1* other influenza virus identified J10.8† G05.1*
Revise non– essential modifiers, delete morphology code	Polycythemia (primary) (rubra) (vera) (acquired) (secondary) NEC (M9950/1) D45 D75.1 - hypoxemic D75.1 - neonatorum P61.1 - nephrogenous D75.1
and revise subterms	- nephrogenous D73.1 - primary D45 - relative D75.1 - rubra vera D45 - secondary D75.1 - vera D45
Revise code:	Polyneuropathy—continued - in—continued
Revise subterms	 malignant neoplasm NEC (M8000/3) (see also Neoplasm, malignant) C80C80† G63.1* Polyneuropathy, (peripheral) (critical illness) in (due to) uremia N188-5 † G63.8*
Add subterms	Polyp, polypous - colon K63.5 - adenomatous (M8210/0) — see Polyp, adenomatous - inflammatory K51.4
Delete morphology code and revise code	Polyposis – see also Polyp – familial (M8220/0) D12.6 – intestinal (adenomatous) (M8220/0) D12.6 – lymphomatous, malignant (M9677/3) C83.89
Revised code at subterm	Polyposis — (see also Polyp) - intestinal (adenomatous) (8220/0) D12.6 - lymphomatous, malignant C83.91
Revise codes	Poor urinary stream R39.1

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	- vision NEC <u>H54.7</u> <u>H54.9</u>
Add lead term	Postpolio syndrome G14
Add subterms	Postpoliomyelitic – see also condition
	– osteopathy M89.6
	<u>– syndrome G14</u>
	Pre-eclampsia O14.9
	- with pre-existing hypertension O11
	- affecting fetus or newborn P00.0
Revise code	- mild Q13 · <u>Q14.0</u>
	- moderate O14.0
	- severe O14.1
Danies ander	- superimposed O11
Revise codes	Pregnancy (single) (uterine)
	- complicated by (see also Pregnancy, management, affected by) early delivery (with spontaneous labor) NEC 060.1
	early derivery (<u>with spontaneous labor</u>) NEC 600.1
	onset of contractions before 37 weeks'gestation Q60- see Labor, early onset
	Pregnancy (single) (uterine)
	- complicated by (see also Pregnancy, management affected by)
	conditions in
	A00-A07, O98.8
	A08, O98.5
	A65-A79,O98.8
Revise code:	A80-B09, O98.8 <u>O98.5</u>
	Pregnancy (single)(uterine)
	- complicated by
Add subterm:	chloasma (gravidarum) O26.8
	cholestasis (intrahepatic) O26.6
	Pregnancy (single) (uterine)
	-complicated by (see also Pregnancy, management, affected by)
D i L	salpingo-oophoritis O23.5
Revise subterms:	septicemia sepsis (conditions in A40, A41) O98.8
	Pregnancy- continued
	- complicated by – continued
	amnionitis O41.1

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Add subterm:	anaphylactoid syndrome of pregnancy O88.1 anemia (conditions in D50-D64) O99.0
	superfetation O30.8
	<u>syndrome</u> anaphylactoid of pregnancy O88.1
	syphilis (conditions in A50-A53) O98.1
Add subterms and	Pregnancy (single)(uterine)
codes	 complicated by (see also Pregnancy, management, affected by) abnormal, abnormality
	placenta, placental (vessel) O43.1
	<u> accreta O43.2</u>
	<u> increta O43.2</u> <u> morbidly adherent O43.2</u>
	morbidiy adherent 045.2 percreta 043.2
	missed
	abortion O02.1 delivery O36.4
	derivery 030.4 morbidly adherent placenta 043.2
	– necrosis, liver (conditions in K72.–) O26.6
	– – neoplasms NEC O99.8
	phlebothrombosis (superficial) O22.2
	deep O22.3
	- – placenta, placental
	abnormality O43.1
	abruptio or ablatio (<i>see also</i> Abruptio placentae) O45.9 accreta O43.2
	dysfunction O43.8
	<u> increta O43.2</u> malposition (with hemorrhage) O44.1
	maiposition (with hemorrhage) 044.1
	morbidly adherent O43.2
	<u>– – – percreta O43.2</u>
	previa (with hemorrhage) O44.1

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	without hemorrhage O44.0
Add subterms	Pregnancy (single)(uterine)
	- complicated by (see also Pregnancy, management, affected by)
	anemia (conditions in D50–D64) O99.0
	conditions in
	B15-B19, O98.4
	B20-B24, O98.7
	R73.0, O99.8
	high head at term O32.4
	– human immunodeficiency virus [HIV] disease O98.7
	 – insufficient weight gain O26.1
	Pregnancy (single) (uterine)
	- complicated by (see also Pregnancy, management affected by)
	postmaturity O48
Revise code	pre-eclampsia O14.9
	mild O13 <u>O14.0</u>
	moderate O14.0
	severe O14.1
Add subterms	premature rupture of membranes — see Rupture, membranes, premature Pregnancy (single) (uterine)
Add subterins	
	- cervical O00.8
	- complicated by (see also Pregnancy, management, affected by)
	incompetent cervix O34.3
	infection(s)
	with a predominantly sexual mode of transmission NEC O98.3
	amniotic fluid or sac O41.1
	bladder O23.1 genital organ or tract O23.5
	genitar organ of tract O23.3 genitourinary tract NEC O23.9
	genicuminaly fractific 025.7

ALPHABETIC INDEX ENTRIES
idney O23.0 with a predominantly sexual mode of transmission NEC O98.3 pecified NEC O98.8
nancy (single) (uterine) nancy (single) (uterine) nplicated by (see also Pregnancy, management, affected by) ema O12.0 vith gestational hypertension, mild (see also Preeclampsia) O13-O14.0
ukemia <u>(related to alkylating agent)</u> (related to Epipodophyllotoxin) (related to therapy) (syndrome) (M9989/1) D46.9
nature — see also condition very (with spontaneous labor) NEC O60.1 thout spontaneous labor (cesarean section) (induction) O60.3
ence (of) temic lupus erythematosus inhibitor D68.6
ciac (iac (ibrillator (functional) Z95.0 (iac (ibrillator (functional) Z95.0 (iac (ibrillator (functional) Z95.0 (iac (iac iac (iac iac iac iac iac iac iac iac iac iac
sure , skin ulcer L89 ge L89.0 L89.1 IL89.2 V L89.3 Posis (chronic) (skin) L89.2
ge

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Add subterms	III L89.2
	<u>IV L89.3</u>
	- sore (chronic) L89 <u></u>
Add subterms	stage
	<u>IL89.0</u> IIL89.1
	II L89.1 III L89.2
	IV L89.3
	- spinal cord G95.2
	- ulcer (chronic) L89 <u></u>
Add subterms	stage
	<u>IL89.0</u>
	<u> II L89.1</u> III L89.2
	IV L89.3
Add lead term	Presyncope R42
	Problem
Revise Code	– personality F61 - <u>F69</u>
Add subterms	Problem (related to) (with)
	- impaired mobility - due to prolonged bedrest R26.3
	- requiring care provider Z74.0
	Problem (related to)(with)
Revise code	cight H54.7 H54.0
Revise code Revise code	- sight <u>H54.7 H54.9</u> Proctitis K62.8
Revise code	- amebic (acute) $A06.08$
Revise code	Proctocolitis, mucosal K51.5 K51.3
	Prolapse, prolapsed
Delete non-	urethra (acquired) (infected) (mucosa) N36.3
essential modifier	- congenital Q64.7
and subterms	female N81.0

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	<u> with</u> cystocele N81.1
	cystoceie 1831.1 prolapse of uterus – see Prolapse, uterus
	<u> male N36.3</u>
Add lead term	Proliferation of primary cutaneous CD30-positive T-cells C86.6
Add subterms	Prolonged
	- labor O63.9
	affecting fetus or newborn P03.8 first stage O63.0
	- second stage O63.1
	- QT interval R94.3
	- uterine contractions in labor O62.4
	affecting fetus or newborn P03.6
	Proteinuria R80
	- Bence Jones NEC R80 - complicating pregnancy, childbirth or puerperium O12.1
	- complicating pregnancy, clindon in or puerperium 012.1 — significant, with gestation hypertension (see also Pre eclampsia) 014.9
5.1	— superimposed on pre existing hypertensive disorder O11
Delete subterm	- gestational O12.1
	with edema O12.2
	- postural N39.2
	— pre eclamptic (see also Pre eclampsia) O14.9 — affecting fetus or newborn P00.0
Add note	Pseudomonas
Tida note	- aeruginosa, as cause of disease classified elsewhere B96.5
	- mallei infection A24.0 - see Infection, Burkholderia, mallei
	as cause of disease classified elsewhere <u>B96.5B96.8</u>
	- pseudomallei, as the cause of disease classified elsewhere B96.8
Add lead term	Pseudotrichinosis M33.1
Add non–essential modifier	Pseudotuberculosis(extra-intestinal) A28.2
	Puerperal, puerperium
Revise code and	- hemiplegia, cerebral O99.4 O99.3
add subterm:	due to cerebral vascular disorder O99.4
	Puerperal, perperium

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Add subterm	- necrosis, liver (acute)(subacute)(conditions in category K72.0) O90.8 - with renal failure O90.4 - neuritis O90.8 - paralysis, bladder (sphincter) O90.8
Add non-essential modifiers	Pulpitis(acute)(chronic)(hyperplastic) (irreversible)(suppurative)(reversible) (ulcerative) K04.0
Delete apostrophe	Purpura D69.2 - hypergammaglobulinemia (benign)(Waldenström's) D89.0
Revise subterms:	Pyelonephritis (see also Nephritis, tubulo-interstitial) N12 - in (due to) sarcoidosis D86.8† N16.2* septicemia sepsis NEC A41† N16.0*
Add subterm	Pyelitis (congenital)(uremic)(see also Pyelonephritis) N12 - acute - with calculus (impacted)(recurrent) N20.9
Revise subterms	Pyelocystitis (see also Pyelonephritis) N12 – with calculus (impacted)(recurrent) N20.9
Delete morphology codes	Pyelonephritis (see also Nephritis, tubulo–interstitial) N12 – in (due to) – leukemia NEC (M9800/3) C95.9† N16.1* – lymphoma NEC (M9590/3) C85.9† N16.1* – multiple myeloma (M9732/3) C90.0† N16.1*
Revise subterms	Pyelonephritis (see also Nephritis, tubulo–interstitial) N12 – calculous NEC N20.9 – with hydronephrosis N13.2
Revise cross reference:	Pyemia, pyemic (purulent) (see also Septicaemia Sepsis) A41.9
Revise cross reference:	Pyosepticemia — see Septicaemia Sepsis
Add subterm	Pyothorax J86.9 - tuberculous NEC A16.5 with bacteriological and histological confirmation A15.6
Add lead term	Rasmussen encephalitis or syndrome G04.8
Revise subterms	Reflux – ureteral N13.7

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	 – with pyelonephritis (chronic) N11.0 – and calculus (impacted)(recurrent) N20.9
Add subterms	Reflux - esophageal K21.9 - neonatal P78.8 - gastroesophageal K21.9 - in newborn P78.8
Add lead term	Refractory cancer (neoplasm) U85
Add subterm	Removal (from) (of) - cardiac pulse generator (battery) (end-of-life) Z45.0 - catheter (indwelling) (urinary) Z46.6 - device - cardiac NEC Z45.0 - contraceptive Z30.5
Add lead term and	Resistance
subterm	- activated protein C [factor V Leiden mutation] D68.5
Add subterms and revise codes	Resistance (by bacterial agent), resistant (to) - antibiotic, antibiotics U83.9 betalactam NOS U82.9 extended spectrum betalactamase (ESBL) U82.2 methicillin U80.4 U82.1 multiple U88 U83.7 penicillin U80.0 U82.0 penicillin related U80.8 quinolones U83.2 specified (single) NEC U89.8 U83.8 multiple U88 U83.7 vancomycin U81.0 U83.0 vancomycin U81.0 U83.1 - antifungal drug(s) U84.1 - antimicrobial drug U84.9 multiple U84.7 - antineoplastic drug(s) U84.0 - antiviral drug(s) U84.0 - antiviral drug(s) U84.2 - tuberculostatic drug(s) U84.3

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Add lead term	Retention, retained—continued
	Reticular erythematous mucinosis L98.5
	Reticulation, dust – see Pneumoconiosis
Delete	Reticuloendotheliosis
morphology	- acute infantile (M9722/3) C96.0
codes, revise	 leukemic (M9941/3) C91.4 malignant (M9712/3) C85.7 C96.9
codes	- nonlipid (M9722/3) C96.0
Delete	Reticulolymphosarcoma (diffuse) (M9675/3) C83.2 C85.9
morphology	 follicular (M9691/3) C82.19 nodular (M9691/3) C82.19
codes, revise	— nodulai (141707173) C82. 1 2
Delete	Reticulosarcoma (diffuse) (M9591/3) C83.3
morphology code	
Delete	Reticulosis (skin)
morphology	acute, of infancy (M9722/3) C96.0hemophagocytic, familial D76.1
codes, revise	- histiocytic medullary (M9720/3) C96.19
Codes	- lipomelanotic I89.8
	- malignant (midline) (M9713/3) C85.7 <u>C86.0</u>
	 nonlipid (M9722/3) C96.0 polymorphic (M9713/3) C85.7 C83.8
	- porymorphic (M97/13/3) C63.7 C63.6 - Sézary's (M9701/3) C84.1
Revise code at	Reticulosis
subterm	– acute, of infancy C96.0
	Retinitis (see also Chorioretinitis) H30.9
Revise codes:	- albuminurica N18. 8 5† H32.8*
A 111 1.	- renal N18.85† H32.8* Richter syndrome C91.1
Add lead term	
	Rokitansky-Aschoff sinuses (gallbladder) K82.8

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Add lead term:	Romano-Ward syndrome I45.8
	Romberg's disease or syndrome G51.8
Add lead term:	Rotation
	- anomalous, incomplete or insufficient, intestine Q43.3
	Rotes Quérol disease or syndrome M48.1
	Roth(-Bernhardt) disease or syndrome (meralgia paraesthetica) G57.1
	Rupture, ruptured
	···
Revise code and	- appendix (with peritonitis) K35.0
add subterms:	<u>with</u>
	peritonitis, generalized K35.2
	peritonitis, localized K35.3
Delete alternative	Rupture, ruptured - vagina S31.4
Defete afternative	- vagina 351.4 - complicating delivery (see also Laceration, vagina, complicating delivery) O71.4
Add asterisk code	Sarcocele (benign)
Add asterisk code	- syphilitic (late) A52.7 †N51.1*
	congenital A50.5 <u>† N51.1*</u>
Delete	Sarcoma (M8800/3) – see also Neoplasm, connective tissue, malignant
morphology	
codes, revise	- dendritic cells (accessory cells) (follicular) (interdigitating)C96.4
codes	- granulocytic <u>(M9930/3)</u> C92.3
	 histiocytic C96.8 Hodgkin's (M9662/3) C81.7 C81.3
	- Hodgkin <u>-s (M9602/3) C81.7 C81.5</u> - immunoblastic (diffuse) (M9684/3) C83.43
	- Infinitioblastic (uffuse) (1415-05-473) C63:43
	- lymphoblastic (M9685/3) C83.5
	- lymphocytic (M9670/3) C83.0 C85.9
	– mast cell (M9740/3) C96.2
	– myeloid <u>(M9930/3)</u> C92.3
	– plasma cell <u>(M9731/3)-C90.23</u>
	- pleomorphic cell (M8802/3)
	- reticulum cell (diffuse) (M9593/3) C83.3
	nodular (M9698/3) C82.2 <u>C96.9</u> pleomorphic cell type (M9680/3) C83.3
	Sarcomatosis
	Sat Colliatosis

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Revise code and add subterms:	unspecified site (M8800/6) C80C79.9 - primary site not indicated C80.9 - primary site unknown, so stated C80.0
	Sarcomatosis
Delete subterms	 unspecified site (M8800/6) C79.9 primary site not indicated C80.9 primary site unknown, so stated C80.0
Delete morphology codes, revise codes	Schilling-type monocytic leukemia (M9890/3)-C93.90
Revise code	Schüller-Christian disease or syndrome D76.0-C96.5
Amend index lead term	Schultz's disease or syndrome D70
Correct typographical error	Schweninger Buzzi anetoderma L90.1
Delete subterm	Sclerosis, sclerotic - renal N26 arteriolar (hyaline)(hyperplastic)(see also Hypertension, kidney)I12 with cystine storage disease E72.0† N29.8* - hypertension (see also Hypertension, kidney) I12
Add lead term	Self neglect R46.8 - causing insufficient intake of food and water R63.6
Revise code	Sepsis - umbilical, umbilical cord (newborn) (unspecified species) P38-P36.— Sepsis (generalised) (see also Septicaemia Infection) A41.9
Revise cross reference and revise subterms:	- actinomycotic A42.7 - adrenal haemorrhage syndrome (meningococcal) A39.1+ E35.1* - anaerobic A41.4 - Bacillus anthracis A22.7 - bacterial, newborn P36.9 - due to

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	anaerobes NEC P36.5 Escherichia coli P36.4
	Escherichia con P36.4 Staphylococcus NEC P36.3
	aureus P36.2
	streptococcus NEC P36.1
	group B P36.0
	specified type NEC P36.8
	- Brucella (see also Brucellosis) A23.9
	- candidal B37.7
	- cryptogenic A41.9
	- due to device, implant or graft (see also Complications, by site and type, infection or inflammation) T85.7
	specified NEC T85.7
	urinary NEC T83.5
	vascular NEC T82.7
	ventricular intracranial shunt T85.7
	- during labour O75.3
	- Erysipelothrix (erysipeloid) (rhusiopathiae) A26.7
	- Escherichia coli A41.5
	- extraintestinal yersiniosis A28.2
	- following
	abortion (subsequent episode) O08.0
	current episode — see Abortion - ectopic or molar pregnancy O08.0
	ectopic of moral pregnancy 608.0
	infusion, therapeutic injection or transfusion T80.2
	- gangrenous A41.9
	- gonococcal A54.8
	- Gram-negative (organism) A41.5
	anaerobic A41.4
	- Haemophilus influenzae A41.3
	- herpesviral B00.7
	- intraocular H44.0
	- Listeria monocytogenes A32.7
	- localised, in operation wound T81.4 (see Infection) - meliodosis A24.1
	- meningeal – see Meningitis

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	- meningococcal A39.4
	<u> acute A39.2</u>
	chronic A39.3 - newborn NEC P36.9
	due to anaerobes NEC P36.5
	Escherichia coli P36.4
	Staphylococcus P36.3
	aureus P36.2
	streptococcus P36.1
	group B P36.0
	specified NEC P36.8
	- Pasteurella multocida A28.0
	- pelvic, puerperal, postpartum, childbirth O85
	- pneumococcal A40.3
	- puerperal, postpartum, childbirth (pelvic) O85
	- Salmonella (arizonae) (cholerae-suis) (enteritidis) (typhimurium) A02.1
	- Shigella (see also Dysentery, bacillary) A03.9
	- specified organism NEC A41.8
	- Staphylococcus, staphylococcal A41.2
	<u> aureus A41.0</u>
	coagulase-negative A41.1
	specified NEC A41.1
	- Streptococcus, streptococcal A40.9
	agalactiae A40.1
	<u> group</u>
	<u> A A40.0</u> <u> B A40.1</u>
	D A40.1 D A40.2
	neonatal P36.1
	- pneumoniae A40.3
	pyogenes A40.0
	specified NEC A40.8
	- tracheostomy stoma J95.0
	- tularemic A21.7
	- umbilical (newborn) (organism unspecified) P38
	tetanus A33

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	- urinary N39.0
	- Yersinia pestis A20.7
	Sepsis (generalized) (see also Septicemia) A41.9 - puerperal, postpartum, childbirth (pelvic) O85
	- Salmonella (arizonae) (cholerae-suis) (enteritidis) (typhimurium) A02.1
	- severe, as a result of disease classified elsewhere R65.1
Add subterm:	- Shigella (<i>see also</i> Dystentery, bacillary) A03.9
	Septicemia, septicemic (generalised) (suppurative)A41.9 see Sepsis
Revise cross	-actinomycotic A42.7
reference and	-adrenal haemorrhage syndrome (meningococcal) A39.1† E35.1*
delete subterms:	-anaerobic A41.4
	Bacillus anthracis A22.7
	-Brucella (see also Brucella) A23.9
	-candidal B37.7
	-cryptogenic A41.9
	-due to infusion, therapeutic injection or transfusion T80.2
	-during labour O75.3
	—Erysipelothrix (erysipeloid) (rhusiopathiae) A26.7 —Escherichia coli A41.51
	- extraintestinal yersiniosis A28.2 -following
	— abortion (subsequent episode) 008.0
	— current episode — see Abortion
	— ectopic or molar pregnancy 008.0
	— immunisation T88.0
	infusion, therapeutic injection or transfusion T80.2
	gangrenous A41.9
	gonococcal A54.8
	Gram negative (organism) A41.5
	— anaerobic A41.4
	Haemophilus influenzae A41.3
	- herpesviral B00.7
	-Listeria monocytogenes A32.7
	-melioidosis A24.1
	-meningeal see Meningitis
	-meningococcal A39.4
	<u>acute A39.2</u>

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	— chronic A39.3
	-newborn NEC (see also Sepsis, newborn) P36.9
	— Pasteurella multocida A28.0
	pneumococcal A40.3
	- postprocedural T81.4
	- puerperal, postpartum O85
	—Salmonella (arizonae) (cholerae suis) (enteritidis) (typhimurium) A02.1
	-Shigella (see also Dysentery, bacillary) A03.9
	-specified organism NEC A41.8
	-Staphylococcus, staphylococcal A41.2
	aureus A41.0
	— coagulase negative A41.1
	— specified NEC A41.1
	-Streptococcus, streptococcal A40.9
	— agalactiae A40.1
	group
	A A40.0
	— B A40.1
	— D A40.2
	— neonatal P36.1
	— pneumoniae A40.3
	— pyogenes A40.0
	specified NEC A40.8
	-tularaemic A21.7
	Yersinia pestis A20.7
	Sequelae
.	- childbirth complication O94
Revise codes:	- resulting in death (one year or more after delivery) O97
	– – between 42 days and one year after delivery O96 <u>.</u> -
	- delivery complication O94
	- resulting in death (one year or more after delivery) O97
	– – between 42 days and one year after delivery O96 <u>.</u> -
	- pregnancy complication(s) O94
	- resulting in death (one year or more after delivery) O97
	– – between 42 days and one year after delivery O96 <u>.</u> -

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	– puerperium complication(s) O94
	 resulting in death (one year or more after delivery) O97 between 42 days and one year after delivery O96
Revise lead term and delete morphology code	Sézary ² s disease, reticulosis or syndrome (M9701/3) C84.1
Revise code	Shock R57.9 -septic A41.9 R57.2
Revise cross reference	Shutdown, renal (see also Failure, renal kidney) N19
Revise code and add subterm Delete non- essential mod	Sigmoiditis (see also Enteritis) A09 A09.9 - infectious A09.0 - noninfectious (presumed) K52.9
Add subterms	Silo-filler's disease J68.8 - bronchitis J68.0 - pneumonitis J68.0 - pulmonary edema J68.1
Revise code	Sinusitis (accessory) (chronic) (hyperplastic) (nasal) (nonpurulent) (purulent) J32.9 – allergic (see also Rhinitis, allergic) J30.3-J30.4
Revise	Sore - chiclero B55.1 - Delhi B55.1 - desert (see also Ulcer, skin) L98.4 - eye H57.1 - Lahore B55.1 - mouth K13.7 - muscle M79.1 Naga (see also Ulcer, skin) L98.4 - oriental B55.1 - pressure L89
Add subterms	stage II L89.0 II L89.1

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	III L89.2 IV L89.3
Add dagger symbol and asterisk code	Splenomegaly, splenomegalia R16.1 – Egyptian B65.1† D77*
Delete code at subterm level	Spots - Bitot's H11.1 - in the young child E50.1† H13.8*
Add lead term	Steatohepatitis (nonalcoholic) K75.8
Add non–essential modifier	Stenosis - organ or site, congenital NEC – see Atresia, by site - pulmonary (artery)(congenital) (supravalvular)(subvalvular) Q25.6
Add subterm:	Strain -see also Sprain - muscle M62.6 - traumatic see Injury, muscle, by site
Revise cross reference:	Streptococcemia (see also Septicaemia-Sepsis, streptococcal) A40.9
Revise code	Stricture (see also Stenosis) R68.8 eustachian tube H68.1 - congenital Q17.8 Q16.4
Revise subterm	Subluxation - see also Dislocation complex-(vertebral) M99.1
Revise code and add subterms	Sudden -death, cause unknown R96.0 infant R95.9 with mention of autopsy R95.0 - infant death syndrome R95.9 with mention of autopsy R95.0
Revise subterm and cross reference	Suppression - renal kidney (see also Failure, renal kidney) N19

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Add lead term:	Swollen – see Swelling Swyer's syndrome Q99.1 Sycosis L73.8
Modify subterm	Syncope R55 - fatal <u>unexplained</u> R96.0
Modify subterm Add subterms	Syndrome – see also Disease - antiocardiolipin D68.8 D68.6 - antiphospholipid D68.6
Add subterms	Syndrome – see also Disease - systemic inflammatory response
	infectious origin with organ failure R65.1 without organ failure R65.0
	non-infectious origin with organ failure R65.3 without organ failure R65.2
Add subterms	without organ faiture R65.2 unspecified origin R65.9 Syndrome — see also Disease
Add subternis	- lobotomy F07.0 - long QT 145.8
	- low
	- prolonged QT 145.8 Syndrome- see also Disease
Add subterm	- amyostatic (Wilson's disease) E83.0 - anaphylactoid, of pregnancy O88.1
ridd subterin	- androgen resistance E34.5 Syndrome - see also Disease
	milk-alkali E83.5
Add index subterm:	- Miller Fisher G61.0 - monofixation H50.4
	Syndrome—continued
Revise code and	 generalized, neoplastic (malignant) C80C79.9 primary site not indicated C80.9
add subterms:	primary site unknown, so stated C80.0

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Revise subterms:	Syndrome – see also Disease - systemic inflammatory response as a result of disease classified elsewhere R65.9 - infectious origin with organ failure (severe sepsis) R65.1 without organ failureR65.0 - non-infectious origin with organ failure R65.3 without organ failure R65.2 - unspecified origin, as a result of disease classified elsewhere R65.9 Syndrome — continued
Add subterm:	trisomy NEC Q92.9 22 Q92.8 - tumor lysis (following antineoplastic drug therapy)(spontaneous) NEC E88.3
Revise cross reference:	- twin (to twin) transfusion Syndrome — see also Disease uremia, chronic (see also Failure, renal Disease, kidney, chronic) N18.9
Add subterm	Syndrome – see also Disease – postphlebitic I87.0 – postpolio (postpoliomyelitic) G14 – postvagotomy K91.1 Syndrome – see also Disease
Add subterm	- depersonalization(–derealization) F48.1 - distal intestinal obstruction E84.1
Add subterm	Syndrome - postthrombotic I87.0
Revise non– essential modifier and add subterm	Syndrome (see also Disease) - adrenocortical (see Cushing's)E27.0 - with Cushing's syndrome -see Cushing's

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Delete morphology codes, revise codes and add non-essential modifiers	Syndrome – see also Disease - 5q-minus (related to alkylating agent) (related to Epipodophyllotoxin) (related to therapy) D46.6 - hemophagocytic, infection-associated D76.2 - histiocytosis NEC D76.3 - myelodysplastic (M9989/1) (related to alkylating agent) (related to Epipodophyllotoxin) (related to therapy) D46.9 - specified NEC D46.7 - myeloproliferative (chronic) (M9960/1) D47.1 - preleukemic (M9989/1) D46.9
Add subterm	Syndrome – see also Disease - nail patella Q87.2 - neonatal abstinence P96.1 - nephritic (see also Nephritis) N05.–
Add subterm	Syndrome – see also Disease - headache NEC G44.8 - HELLP (hemolysis, elevated liver enzymes and low platelet count) O14.2 - haemolytic-uremic D59.3
Add subterms	Syndrome — see also Disease - coronary, intermediate I20.0 - intermediate I20.0 - slow flow I20.8 - sirenomelia Q87.2 - slow flow, coronary I20.8 - spasmodic - upward movement, eyes H51.8 - winking F95.8
Add subterm	Syndrome- continued cat-cry Q93.4 - cat eye Q92.8
Revise code and add subterm	Syndrome - see also Disease

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	- sudden infant death R95.9
	with mention of autopsy R95.0 Syndrome—continued
Delete subterms	– generalized, neoplastic (malignant) C79.9
	— primary site not indicated C80.9
	— primary site unknown, so stated C80.0
	Synovitis—continued – in (due to)
Revise code	- in (due to) crystals M11.9-M65.8
Revise code	System, systematic – see also condition
110 1100 0000	- lupus erythematosus M32.9
	inhibitor present D68.8 D68.6
Add subterms	System, systemic – see also condition
	- atrophy - multiple (brain)(CNS) G90.3
	- Indusple (brain)(CNS) G90.5 - disease, combined – see Degeneration, combined
	- lupus erythematosus M32.9
	inhibitor present D68.86
Add subterms and	Tachycardia R00.0
non-essential	- atrial I47.1
modifier	- atrioventricular (re-entrant) I47.1
	- auricular I47.1 - fetal - see Distress, fetal
	- junctional I47.1
	- nodal I47.1
	- paroxysmal <u>NEC</u> 147.9
	atrial I47.1
	atrioventricular (AV) (re-entrant) I47.1
	junctional I47.1 nodal I47.1
	nodar 147.1 supraventricular I47.1
	- ventricular I47.2
Revise code and	Tag (hypertrophied skin) (infected) L91.8
subterm	
	- anus <u>K64.4I84.6</u>
	- hemorrhoidal <u>K64.4 184.6</u>

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	 - rectum <u>K64.4I84.6</u>
Delete subterm	Tag
Revise codes and add subterm	rectum K64.4 Tear, torn (traumatic) - see also Wound, open - periurethral tissue, obstetric trauma O71.5 O70.0 urethra, obstetric trauma O71.5 - periurethral tissue O70.0 - uterus - see Injury, uterus
Delete subterm	Tear, torn (traumatic) (see also Wound, open) T14.1 urethra, obstetric trauma O71.5 — periurethral tissue O70.0
Add subterms	Telangiectasia, telangiectasis (verrucous) I78.1 - ataxic (cerebellar) (Louis-Bar) G11.3 - hemorrhagic, hereditary (congenital) (senile) I78.0 - juxtafoveal H35.0 - macular H35.0 - parafoveal H35.0 - retinal (idiopathic)(juxtafoveal)(macular) (parafoveal) H35.0 - spider I78.1
Revise code	Tenosynovitis M65.9 - in (due to) - crystals M11.9 M65.8
Revise and add subterm	Test(s) blood-drug Z04.0 - cardiac pulse generator (battery) Z45.0 - device (implanted) NEC Z45.0 - pulse generator (battery) Z45.0
Delete morphology codes	Thrombocythemia (essential) (hemorrhagic) (idiopathic) (primary) (M9962/1) D47.3 Thrombocytosis, assential, D75.2 D47.3
Revise code Add lead term and	Thrombocytosis, essential D75.2 D47.3 Thrombophilia

INSTRUCTION	ALPHABETIC INDEX ENTRIES
subterms	- primary D68.5
Revise code:	Thrombosis, thrombotic (multiple) (progressive) (septic) (vein) (vessel) I82.9 - appendix, septic <u>K35.9 K35.8</u>
Revise code and subterm	Thrombosis, thrombotic (multiple) (progressive) (septic) (vein) (vessel) I82.9 - anal (see also Thrombosis, perianal venous) I84.3K64.5 - penis, penile N48.8 - perianal I84.3venous K64.5 - peripheral arteries I74.4
	- vena cava (inferior) (superior) I82.2 <u>- venous, perianal K64.5</u>
Revise cross reference:	Toxemia R68.8 -bacterial — see Septicaemia Sepsis - pre-eclamptic (see also Pre-eclampsia) O14.9 - septic (see also Septicaemia Sepsis) A41.9
Add subterm	Tracheitis (acute) (catarrhal) (infantile) (membranous) (plastic) (pneumococcal) (viral) J04.1 – with – bronchitis (15 years of age and above) J40 – - tuberculous NEC A16.4 – with bacteriological and histological confirmation A15.5
Add non-essential modifier and code:	Transfusion - blood (session) Z51.3 - reaction or complication -see Complications, transfusion - without reported diagnosis Z51.3
Add subterm:	Transplant(ed)(status) Z94.9 pancreas Z94.8 - skin Z94.5 - social Z60.3 -specified organ or tissue NEC Z94.8 - stem cells Z94.8
Amend index as follows	Treponematosis due to - due to - Treponema pallidum (see Syphilis) - Treponema pertenue (see Yaws)

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Revise code	Tuberculosis, tubercular, tuberculous (caseous) (degeneration) (gangrene) (necrosis) A16.9 - abscess NEC (respiratory) A16.9 - dura (mater) (cerebral) (spinal) A17.8† G01* G07*
Revise code	Tuberculosis, tubercular, tuberculous (caseous)(degeneration)(gangrene)(necrosis) A16.9 – heart A18.8† 143.0*152.0*
Revise subterms:	Tularemia A21.9 - pneumonia (any), pneumonic A21.2† J17.0* - septicemia sepsis A21.7 - specified NEC A21.8
Revise codes:	Tumor(M8000/1)-see Neoplasm, uncertain behavior – germ cell (M9064/3) – see also Neoplasm, malignant – mixed (M9085/3) C80C80
	– malignant (M8000/3) – see also Neoplasm, malignant – fusiform cell (type) (M8004/3) C80C80
	- giant cell (type) (M8003/3) C80C80 - mixed NEC (M8940/3) C80C80 - small cell (type) (M8002/3) C80C80
	spindle cell (type) (M8004/3) C80C80 - unclassified (M8000/3) C80C80 Tumour- continued
Add subterms:	- sternomastoid (congenital) Q68.0 - stromal - gastrointestinal (GIST) D37.9 - benign – see Neoplasm, benign - colon D37.4
	esophagus D37.7 - malignant — see Neoplasm, malignant - peritoneum D48.4 - rectum D37.5 - small intestine D37.2
	- <u>smart intestine D37.2</u> - <u>specified site NEC – see Neoplasm, uncertain behaviour</u> - <u>stomach D37.1</u> - sweat gland (M8400/1)- <i>see also</i> Tumor-(M8000/1) – <i>see also</i> Neoplasm, uncertain behaviour
	1 unioi – (1/10000/1) – see uiso i veopiasii, uncertain benavioui

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	- odontogenic (M9270/1) D48.0
	calcifying epithelial (M9340/0) D16.5
Add subterms	upper jaw (bone) D16.4 keratocystic (M9270/0) D16.5
	<u>keratocystic (W19270/0) D16.5</u> upper jaw (bone) D16.4
Delete	Tumor (M8000/1) – see also Neoplasm
morphology	
codes, revise	<u>– Burkitt's (M9687/3)</u> C83.7
codes	- malignant (M8000/3) - see also Neoplasm, malignant
	— mast cell C96.2
	— plasma cell, localized NEC C90.3
	<u>- mast cell (M9740/1) D47.0</u> - malignant (M9740/3) C96.2
	manghant (M97403) C90.2 plasma cell (malignant) (M9731/3) C90.2 C90.3
	Tumor (see also Neoplasm, uncertain behavior) (M8000/1)
	- malignant (M8000/3) - see also Neoplasm, malignant
	mast cell C96.2
Delete one indent	plasma cell, localized NEC C90.3
at subterm	- mast cell D47.0
	malignant C96.2
	— plasma cell (malignant) C90.3
Revise code	Tumor - Klatskin (M8162/3) C22.1 <u>C24.0</u>
Revise	Ulcer, ulcerated, ulcerating, ulceration, ulcerative L98.4
	- cystic duct K82.8
A 11 1.	- decubitus (skin, any site) L89 <u>.</u> -
Add subterms	<u> stage</u>
	<u>IL89.0</u>
	<u> II L89.1</u>
	<u> III L89.2</u>
	- <u>IV L89.3</u>
	cervix N86
	- dendritic, cornea (herpes simplex) B00.5† H19.1* - dysenteric NEC A09 A09.9
	- uyschiche Nee Aub <u>Aub.b</u>

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	- phlebitis — see Phlebitis
Add subterms	- plaster L89 <u></u>
	<u> stage</u>
	<u>IL89.0</u>
	<u> II L89.1</u>
	<u> III L89.2</u> IV L89.3
	- popliteal space — see Ulcer, lower limb
	- postpyloric — see Ulcer, duodenum
	- prepuce N48.1
	- prepyloric — see Ulcer, stomach
Add subterms	- pressure L89 <u></u> stage
	stage I L89.0
	II L89.1
	III L89.2
	IV L89.3
	- pyloric — see Ulcer, stomach
	- rectosigmoid K63.3
	- rectum (solitary) (sphincter) K62.6 - stercoraceous, stercoral K62.6
	varicose — see Varicose, ulcer, anus
	- retina H30.0
	- rodent (M8090/3) — see also Neoplasm, skin, malignant
	- sclera H15.0
	- scrofulous (tuberculous) A18.2 - scrotum N50.8
	tuberculous A18.1† N51.8*
	varicose I86.1
	- seminal vesicle N50.8
	- sigmoid (see also Ulcer, intestine) K63.3
	- skin (atrophic) (chronic) (neurogenic) (perforating) (pyogenic) (trophic) L98.4 - with gangrene (see also Gangrene) R02
	- amoebic A06.7

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	decubitus L89 <u>.</u> -
A 11 1.	<u> stage</u>
Add subterms	<u>IL89.0</u>
	<u>II L89.1</u>
	III L89.2
	IV L89.3 lower limb L97
	tuberculous (primary) A18.4
	- varicose — see Ulcer, varicose
	- sloughing NEC — see Ulcer, skin
	Ulcer, ulcerated, ulcerating, ulcerative L98.4
	- foot (indolent) (see also Ulcer, lower limb) L97
Add non-essential modifier and	varicose (venous) 183.0 inflamed or infected 183.2
subterms:	Inflatfied of fiffected 183.2
Subtering.	venous NEC 183.0
	due to venous insufficiency I87.2
	inflamed or infected I83.2
	<u> postphlebitic (infected) (inflamed) (postthrombotic) I87.0</u>
	- lower limb (atrophic) (chronic) (neurogenic) (perforating) (pyogenic) (trophic) (tropical) L97
	varicose (venous) I83.0
	inflamed or infected I83.2
	<u> venous NEC I83.0</u>
	due to venous insufficiency I87.2
	inflamed or infected I83.2
	postphlebitic (infected) (inflamed) (postthrombotic) I87.0
	- stasis (venous) I83.0
	- inflamed or infected I83.2
	variance (lawer limb, any part) (vanous) IV2 0
	- varicose (lower limb, any part) (venous) I83.0 - inflamed or infected I83.2

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	- venous NEC I83.0
	due to venous insufficiency I87.2
	inflamed or infected I83.2
	postphlebitic (infected) (inflamed) (postthrombotic) I87.0 Ulcer, ulcerated, ulceration, ulcerative L98.4
	· · · · · · · · · · · · · · · · · · ·
Delete and add	- anus (sphincter) (solitary) K62.6
subterms	- varicose see Varicose, ulcer, anus
	- heel (see also Ulcer, lower limb) L97
	- hemorrhoid (see also Hemorrhoids, by degree) K64.8
	- Hunner's N30.1
	rectum (sphincter) (solitary) K62.6
	stercoraceous, stercoral K62.6
	- varicose see Varicose, ulcer, anus
	- varicose (lower limb, any part) I83.0
	anus see Varicose, ulcer, anus
	esophagus I85.9 bleeding I85.0
	inflamed or infected I83.2
	nasal septum I86.8
	<u>rectum</u> see Varicose, ulcer, anus
Revise cross	Uremia, uraemic (coma) N19
reference:	- chronic (see also Failure, renal Disease, kidney, chronic N18.9
Add subterm	Urethrocele
	- congenital Q64.7
	- female N81.0 with
	with cystocele N81.1
	prolapse of uterus — see Prolapse, uterus
	- male N36.3
	Varicose
	- phlebitis — see Varicose vein, inflamed
Add non-essential	- ulcer (lower limb, any part) (venous) I83.0
modifier	inflamed or infected I83.2

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	- vein (lower limb) (ruptured) I83.9
	inflamed or infected I83.1
	with ulcer (venous) I83.2
	ulcerated I83.0 inflamed or infected I83.2
D 1 . 1.	Varicose
Delete subterms	Varicose
	- ulcer (lower limb, any part) I83.0
	— anus 184.8
	— external I84.4
	— internal I84.1
	esophagus I85.9
	bleeding I85.0
	inflamed or infected I83.2
	nasal septum I86.8
	<u>rectum</u> see Varicose, ulcer, anus
	scrotum I86.1
	- vein (lower limb) (ruptured) I83.9
	- anus see Hemorrhoids
	congenital (peripheral) Q27.8
	rectum see Hemorrhoids, internal
	scrotum (ulcerated) I86.1
Add subterms	Varix (lower limb)(ruptured) I83.9
	- esophagus (ulcerated) I85.9
	bleeding I85.0
	congenital Q27.8
	in (due to)
	alcoholic liver disease K70† I98.2*
	with bleeding K70† 198.3* cirrhosis of liver K74† 198.2*
	Cirrilosis of fiver K74† 198.2* with bleeding K74† 198.3*
	schistosomiasis B65† 198.2*
	with bleeding B65† I98.3*
	toxic liver disease K71† I98.2*
	<u> with bleeding K71† I98.3*</u>

INSTRUCTION	ALPHABETIC INDEX ENTRIES
Add non-essential modifier:	Varix (lower limb) (ruptured) I83.9 - with - inflammation or infection I83.1 with ulcer (venous) I83.2 - stasis dermatitis I83.1 with ulcer I83.2 - ulcer (venous) I83.0 with inflammation or infection I83.2 - inflamed or infected I83.1
	- ulcerated I83.0 - inflamed or infected I83.2
Delete subterms	Varix (lower limb) (ruptured) I83.9anus - see Hemorrhoids - bladder I86.2rectum - see Hemorrhoids, internal - renal papilla I86.8
Add subterm	Vestibulitis (ear) H83.0 – nose (external) J34.8 – vulvar N76.2
Add lead term	Vibrio vulnificus, as cause of disease classified elsewhere B98.1
Revise code	Vincent's - stomatitis A69.0-A69.1
Revise subterms and codes	Vision, visual - binocular, suppression H53.3 - blurred, blurring H53.8 - hysterical F44.6 - defect, defective NEC H54.7H54.9
	- halos H53.1 - loss <u>H54.7H54.9</u> both eyes <u>H54.3H54.1</u>

INSTRUCTION	ALPHABETIC INDEX ENTRIES
	complete, with or without light perception— see Blindness
	one eye <u>H54.6H54.5</u>
	sudden H53.1
	- low (both eyes) H54.2
	one eye (other eye normal) <u>H54.5H54.6</u> blindness, other eye <u>H54.1H54.4</u>
	officiess, other eye fi34.1134.4 -perception, simultaneous without fusion H53.3
D 1 1	Waldenström's
Revise lead term and delete	- hypergammaglobulinemia D89.0
	- syndrome or macroglobulinemia (M9761/3) C88.0
morphology code	Wegener's granulomatosis or syndrome M31.3
	- with lung involvement M31.3 ⁺ J99.1 [*]
Add subterm:	kidney involvement M31.3† N08.5*
Add subterm.	lung involvement M31.3† J99.1*
Add index lead	Werner's disease or syndrome E34.8
term	Werner–Schultz disease D70
Add cross	Wernicke-Korsakov syndrome – see also Korsakov's disease, psychosis or syndrome (alcoholic) F10.6
reference:	
Delete subterm	Xanthoma(s), xanthomatosis (primary)(familial)(hereditary) E75.5
	- bone (generalista) D76.0
Add subterm	Xanthoma(s), xanthomatosis (primary)(familial)(hereditary) E75.5
	<u>– bone (generalisata) C96.5</u>

External causes of injury

INSTRUCTION	EXTERNAL CAUSES OF INJURY INDEX
	Earthquake (any injury) X34.9_
Revise code at lead	- cataclysmic earth movements X34.0
term and add	- specified effect NEC X34.8
subterms:	<u>- tsunami X34.1</u>
Add subterms and	Fall, falling- continued
codes	-in, on
	aircraft NEC V97.0

	with accident to aircraft V97.0
	while boarding or alighting V97.1
	bath(tub) W18
	escalator
	transport vehicle after collision - see Accident, transport, by type of vehicle, collision
	tub W18
	-into
Add subterms	Fall, falling (accidental)
Add subterms	
	- from, off
	cherry picker W17
	lifting device NEC W17
	mobile elevated work platform [MEWP] W17
	sky lift W17
Revise code at lead	Fracture (circumstances unknown or unspecified) X59.90
	Fracture (circumstances unknown of unspective) A57.790
term	
Revise code	Struck (accidentally) by
Revise code	Struck (accidentally) by
	••••
	- object NEC W22
	- object NEC W22 - falling W20
	- object NEC W22 - falling W20 from, in, on
	- object NEC W22 falling W20 from, in, on
Revise code	
	- object NEC W22 - falling W20 from, in, on earthquake X34 Trapped (accidentally)
Revise code	- object NEC W22 - falling W20 from, in, on earthquake X34 Trapped (accidentally) - between
Revise code	- object NEC W22 - falling W20 from, in, on earthquake X34 Trapped (accidentally) - between - buildings (collapsing) in earthquake X34.0
Revise code Add subterms:	- object NEC W22 - falling W20 from, in, on earthquake X34 Trapped (accidentally) - between - buildings (collapsing) in earthquake X34.0 - objects (moving) (stationary and moving) (see also Caught) W23
Revise code Add subterms: Add lead term:	- object NEC W22 - falling W20 from, in, on earthquake X34 Trapped (accidentally) - between - buildings (collapsing) in earthquake X34.0 - objects (moving) (stationary and moving) (see also Caught) W23 Tsunami (any injury) NEC X34.1
Add subterms: Add lead term: Revise code and	- object NEC W22 - falling W20 from, in, on earthquake X34 Trapped (accidentally) - between - buildings (collapsing) in earthquake X34.0 - objects (moving) (stationary and moving) (see also Caught) W23 Tsunami (any injury) NEC X34.1 Victim (of)
Revise code Add subterms: Add lead term:	- object NEC W22 - falling W20 from, in, on earthquake X34 Trapped (accidentally) - between - buildings (collapsing) in earthquake X34.0 - objects (moving) (stationary and moving) (see also Caught) W23 Tsunami (any injury) NEC X34.1 Victim (of) - avalanche X36
Add subterms: Add lead term: Revise code and	- object NEC W22 - falling W20 from, in, on earthquake X34 Trapped (accidentally) - between - buildings (collapsing) in earthquake X34.0 - objects (moving) (stationary and moving) (see also Caught) W23 Tsunami (any injury) NEC X34.1 Victim (of) - avalanche X36 earth movement NEC X36
Add subterms: Add lead term: Revise code and	- object NEC W22 - falling W20 from, in, on earthquake X34 Trapped (accidentally) - between - buildings (collapsing) in earthquake X34.0 - objects (moving) (stationary and moving) (see also Caught) W23 Tsunami (anv injury) NEC X34.1 Victim (of) - avalanche X36 earth movement NEC X36 earthquake X34 X34.9
Add subterms: Add lead term: Revise code and	- object NEC W22 - falling W20 from, in, on earthquake X34 Trapped (accidentally) - between - buildings (collapsing) in earthquake X34.0 - objects (moving) (stationary and moving) (see also Caught) W23 Tsunami (anv injury) NEC X34.1 Victim (of) - avalanche X36 earth movement NEC X36 earthquake X34 X34.9 - cataclysmic earth movements X34.0
Revise code Add subterms: Add lead term: Revise code and	- object NEC W22 - falling W20 from, in, on earthquake X34 Trapped (accidentally) - between - buildings (collapsing) in earthquake X34.0 - objects (moving) (stationary and moving) (see also Caught) W23 Tsunami (anv injury) NEC X34.1 Victim (of) - avalanche X36 earth movement NEC X36 earthquake X34 X34.9

Table of Drugs and Chemicals

INSTRUCTION	ALPHABETIC INDEX ENTRIES									
Revise code:	Antianxiety drug NEC	T43.5	X41	X61	Y11	Y49.5 <u>Y47.9</u>				
Revise table of drugs and chemicals	Cyclobenzaprine	T43.0 T48.1	X41- <u>X44</u>	X61 <u>X64</u>	¥11 - <u>¥14</u>	¥49.0 Y55.1				
Revise table	Helium (nonmedicinal) NEC - medicinal	T59.8 T48.7	<u>X47.</u> _ X 44.	X67. -X64.	<u>Y17.</u> <u>-</u> ¥14.	¥55.7 Y55.7				
Add inday ontay	Mathalanadaisalana	T49.0	X44 X44	X64	<u>Y14</u> Y14	Y56.0				
Add index entry	Methylprednisolone - intravenous	T38.0	X44 X44	X64	Y14	Y42.0				
Revise codes	Phenazopyridine	T37.8 <u>T39.8</u>	X44. X40	X64. <u>X60</u>	Y14. <u>Y10</u>	Y41.8 <u>Y45.9</u>				
Add entries to index	Sildenafil	<u>T46.7</u>	<u>X44</u>	<u>X64</u>	<u>Y14</u>	<u>Y52.7</u>				
Revise codes	Tramadol	T39.3 T40.4	X40. X42	X60. X62	¥10. ¥12	¥45.3 ¥45.0				
Add entries to index	<u>Tadalafil</u>	<u>T46.7</u>	<u>X44</u>	<u>X64</u>	<u>Y14</u>	<u>Y52.7</u>				
Revise index entry in Table of Drugs and Chemicals:	Thallium (compounds) (dust) NEC	T60.4 - <u>T56.8</u>	X48. <u>X49.–</u>	X68 <u>X69</u>	¥18 <u>¥19</u>					
Correct codes as follows	Trichlorethylene	T41.0 T53.2	X44 -X46	X64 <u>X66</u>	¥14- <u>¥16</u>	¥48				

INSTRUCTION	ALPHABETIC INDEX ENTRIES								
Correct codes as follows	Trichloroethane	T53.2	<u>X46</u>	X66	Y16				
Correct codes as follows	Trichloroethylene - vapor NEC	T41.0 T53.2 T53.2 T41.0	X44- X46 X46- X44	X64 X66 X66 X64	¥14- <u>¥16</u> ¥16- <u>¥14</u>	¥48			
Add index entry:	Vardenafil	<u>T46.7</u>	<u>X44</u>	<u>X64</u>	<u>Y14</u>	<u>Y52.7</u>			
Delete index entries	Warfarin - rodenticide - sodium	T45.5 T60.4 X48.	X44.– X48.– X68.	X64 X68	Y14 Y18 Y18.	Y44.2			