

**ERRATUM ITEM LIST**

**HP05-2024DI**

Item No	Item Specification	UNIT (Use for Estimate & Price)	Estimate
1	Barium sulfate 4.6% m/m suspension for CT scan examinations, 225ml	Each	6,310
2	Barium sulfate 96% powder for suspension, for rectal use. Prefilled enema kit with disposable tubing and tip, 397g	Each	2,140
3	Barium sulfate 98% m/m powder for suspension for oral use, 340g	Each	16,173
4	Fluorescein strips 1mg, 100 strips	1 Pack of 100 strips	72
5	Fluorescein 100mg/ml, injection, 5ml	Each	805
6	Gadobenic acid 604.72mg/ml, 7.5ml prefilled syringe	Each	3,050
7	Gadopentetic Acid 469.1mg/ml injection, 20ml	Each	18,640
8	Iohexol 300mg iodine/ml injection, 100ml	Each	374,700
9	Iohexol 300mg iodine/ml injection, 20ml	Each	8,490
10	Iohexol 300mg iodine/ml injection, 50ml	Each	135,400
11	Iohexol 350mg iodine/ml injection, 100ml	Each	355,480
12	Iohexol 350mg iodine/ml injection, 50ml	Each	88,700
13	Iopamidol 300mg iodine/ml injection, 50 ml	Each	750
14	Iopamidol 300mg iodine/ml injection, 100 ml	Each	1,500
15	Iopamidol 370mg iodine/ml injection, 100ml	Each	6,716
16	Iopamidol 370mg iodine/ml injection, 50ml	Each	2,500
17	Iopromide 300mg iodine/ml injection, 100ml	Each	2,188
18	Iopromide 300mg iodine/ml injection, 50ml	Each	21,000
19	Iopromide 370mg iodine/ml injection, 100ml	Each	6,177
20	Meglumine amidotrizoate 66g and sodium amidotrizoate 10g solution, 100ml	Each	35,850
21	Meglumine diatrizoate/amidotrizoate 260mg and sodium diatrizoate/amidotrizoate 40mg injection, 250ml	Each	36,026
22	Meglumine diatrizoate/amidotrizoate 520mg and sodium diatrizoate/amidotrizoate 80mg injection, 250ml	Each	480
23	Lipoarabinoamannan (LAM) antigen urine test strips; 25's; including pipettes; 25's	1 Pack of 25 strips	62,899
24	TUBERCULIN PPD, LIQUID, TEST, 2TU/0.1ML; UNIT FOR INTRADERMAL TESTING BY MANTOUX METHOD 1,5ML VIAL KIT TO INCLUDE TRANSPARENT RULER FOR MEASURING INDURATIONS	Each	237,909
25	URINE TEST STRIPS for the determination of 10 different URINE PARAMETERS i.e. glucose, protein, pH, nitrite, ketone, urobilinogen, bilirubin, blood, leukocytes and specific gravity; 100 strips per container. Strip must have a long handle, facing downwards to be safe and hygienic. Test strip must not be influenced by ascorbic acid. Must be a rapid test with results obtained within 3 minutes. Interpretation of test results must be incorporated on the outside of the immediate container design.	1 Pack of 100 strips	575,022