



health

Department:
Health
REPUBLIC OF SOUTH AFRICA



Private Bag X828, PRETORIA, 0001 Dr AB Xuma Building 1112 Voortrekker Road, Pretoria Townlands 351-JR,
PRETORIA, 0187 Tel (012) 395 8000, Fax (012) 395 8918

Mr E van Zyl
Equity Pharmaceuticals (Pty) Ltd
100 Sovereign Road
Route 21 Corporate Park
Nellmapius Drive
Irene
Pretoria

Dear Mr van Zyl

Section 21 Authorization for ETHAMBUTOL 400MG TABLETS 120'S

Attached, please find the Authorization for exemption under Section 21 of the Medicines and Related Substances Act by SAHPRA granted for:

- **Ethambutol 400mg Tablets 120's**

The quantities for which approval was granted are only estimates based on procurement by provinces over the last 6 months. Please note that the National Department of Health (NDOH) cannot guarantee the procurement of these quantities, as NDOH has no control over orders being placed by provincial depots, and current stock holding might influence estimated quantities.

The following process will be followed to ensure the quality of the product being brought in:

1. Manufacturer will submit an assay and identification of every batch imported.
2. An additional assay of every batch will be done by a quality control laboratory.
3. A random sample will be assayed during the authorized period by a quality control laboratory.
4. Aggregate statistics to be submitted to NDOH in the first week of each month of all orders received and quantities supplied per province.
5. The NDOH needs to be advised of the quantities and date of arrival of stocks in terms of this authorization within 7 days after arrival.
6. The supplier will provide monthly reports, by the 7th of each month, using the attached format of orders received and issues done.
7. Participating Authorities (PAs) will provide a consolidated close out report of usage using the attached format on the date when an authorization lapses.

Section 21 Authorisation re Ethambutol 400mg Tablets 120's 19062026-1


8. The full quantities imported in terms of this Section 21 authorisation must be accounted for.
9. Note that this authorization DOES NOT cover supplies to the private sector.
10. Where this authorization is obtained to provide security of supply due to supply challenges from the contracted supplier, PAs are requested to buy out against contracted suppliers and ensure that related orders are cancelled accordingly to prevent overstocking once the contracted supplier gets back into stock.

It should be noted this authorization applies only for use of the product in the public sector with estimated usage quantities for a period of one month. The authorization is expected to expire on **14 December 2026**.

Table 1: Provincial estimates

Province	Six Months Estimate
DCS	0
EC-MT	5 000
EC-PE	2 000
FS	400
GP	900
KZN	1 200
LP	750
MP	144
NC	300
NW	410
SAMHS	22
WC	800
Total	11 926

Yours sincerely



KHADIJA JAMALOODIEN
CHIEF DIRECTOR: HEALTH PRODUCTS PROCUREMENT

DATE: 19/01/2026

Section 21 Outcome Letter

2026-06-14

Ms Khadija Jamaloodien

1112 Voortrekker Road

Pretoria

buhle.mbongo@health.gov.za

Dear Ms Khadija Jamaloodien

**REQUEST TO USE UNREGISTERED MEDICINE IN TERMS OF SECTION 21 OF THE
MEDICINES AND CONTROLLED SUBSTANCES ACT, 1965 (ACT 101 of 1965):**

Your application dated **2026-06-03** refers

- A. STATUS: Approved**
 - B. APPLICANT: Ms Khadija Jamaloodien**
 - C. IMPORTING COMPANY: EQUITY PHARMACEUTICAL (PTY) LTD**
 - D. NUMBER OF PATIENT/(S) INTENDED TO BE TREATED: 100**
 - E. UNREGISTERED MEDICINES: GENERIC NAME: No Data**
 - F. TRADE NAME: Ethambutol Hydrochloride 400 mg**
 - G. QUANTITY: 11936 Packs (56)**
-
-

H. LETTER NUMBER: S2100030566

Section 21 authorization letters are valid for a period of 6 months from the letter date, unless otherwise specified.


A progress report must be submitted once treatment is completed or on a reauthorization request

Comments:

Yours faithfully,

Dr Shyamli Munbodh

Manager: Section 21 Category A Medicines

A rectangular box containing a handwritten signature in black ink. The signature appears to be 'Shyamli Munbodh' written in a cursive style.

Ms Mahlodi Moropa

Final Approver

S2100030566

A rectangular box containing a handwritten signature in black ink. The signature is stylized and appears to be 'M. P. P.' or similar.




health

Department:
Health
REPUBLIC OF SOUTH AFRICA



Private Bag X828, PRETORIA, 0001 Dr AB Xuma Building 1112 Voortrekker Road, Pretoria Townlands 351-JR, PRETORIA,
0187 Tel (012) 395 8000, Fax (012) 395 8918

REQUEST FOR QUOTATION FORM					
<ul style="list-style-type: none"> Instruction to complete this Request for Quotation (RFQ) PLEASE PROVIDE A QUOTE FOR THE FOLLOWING PRODUCT(S). PLEASE QUOTE ON THIS RFQ FORM AND ATTACH YOUR QUOTE WITH THE REQUESTED DETAILS. THE SECTIONS HIGHLIGHTED IN YELLOW MUST BE COMPLETED BY THE SUPPLIER. THIS DOES NOT CONSTITUTE ANY OBLIGATION TO PROCURE THE ITEM AS THIS WILL BE SUBMITTED FOR CONSIDERATION TO PROVINCIAL PROCUREMENT UNITS TO SERVE AS A BUY OUT AGAINST CURRENT NON-COMPLIANT SUPPLIERS. 					
ONLY RESPONSES FROM DULY REGISTERED SUPPLIERS WILL BE EVALUATED					
REFERENCE NUMBER:	NORMAL		SECTION 21	X	S21RFQ177/2026
QUOTE ENQUIRY DATE	19/05/2026	QUOTE CLOSING DATE		29/05/2026	
FOR CRITICAL DELIVERY, DELIVERY REQUESTED ON/BEFORE (SCM Practitioner to Specify if applicable)					
REQUESTING INSTITUTION CONTACT DETAILS					
NAME OF REQUESTOR	Anzelde Willemse				
EMAIL ADDRESS	anzelde.willemse@health.gov.za				
PHONE No.	066 261 4234	FAX No.	N/A		
PRODUCT INFORMATION					
DESCRIPTION PER MPC	Ethambutol; 400mg; Tablet; 28 Tablets				
TRADE DESCRIPTION	Combutor (Ethambutol) 400mg tablets				
UNIT OF MEASURE	Packs	PACK or BOX (SIZE/ QUANTITY)		120 Tablets	
QUANTITY REQUIRED	11 926				
TO BE COMPLETED BY THE SUPPLIER/ SERVICE PROVIDER					
SUPPLIER CONTACT DETAILS (as per CSD)					
DID YOU BID ITEM FOR TENDER	YES		NO	X	IF NO, WHY Section 21
COMPANY NAME	Equity Pharmaceuticals				
SUPPLIER NUMBER	MAAA007480				
SECURITY CODE					
SUPPLIER CODE (NDoH)					
CONTACT PERSON 1	NAME	Ehrard van Zyl			
	PHONE	012 345 1747	FAX	0123451412	
	MOBILE	072 040 8511			
	E-MAIL	ehrd@equitypharma.co.za			
CONTACT PERSON 2	NAME	Hannes Strydom			
	PHONE	012 345 1747			

	MOBILE	082 616 1954	
	E-MAIL	hannes@equitypharma.co.za	
QUOTE DETAILS			
PRICE PER UNIT (INCL. VAT)	R 106.95	TOTAL PRICE (INCL. DELIVERY & VAT)	R 297 534.90
VOLUMES AVAILABLE – 14DAYS	2782 packs of 120's = 333 840 tablets		
VOLUMES AVAILABLE – 21DAYS			
VOLUMES AVAILABLE – 28DAYS			
VOLUMES AVAILABLE – 35DAYS			
VOLUMES AVAILABLE – 42DAYS			
VOLUMES AVAILABLE – 49DAYS			
VOLUMES AVAILABLE – 56DAYS			
VOLUMES AVAILABLE – 112DAYS			
QUOTE VALIDITY PERIOD NORMAL LEAD/DELIVERY TIME			
DEVIATION TO SPECIFICATION			
<i>COMMENTS: Quote requested for packs of 28's. As per quote, Equity Pharmaceuticals can supply packs of 120's</i>			
Quote requested for 11 926 packs of 28's = 333 928 tablets			
Quoted for 2782 packs of 120's = 333 840 tablets			
DECLARATION BY SUPPLIER			
I hereby declare that in submitting this bid, there has been no consultation, communication, agreement or arrangement with any competitor/supplier regarding the price, quality, quantity, specifications and conditions or delivery particulars of the products or services to which this bid invitation relates.			
NAME	Ehrard van Zyl		
CAPACITY	Business Unit Manager: Specialist Medicine		
SIGNATURE (OF A DULY AUTHORISED REPRESENTATIVE OF THE SUPPLIER)			
DATE	29/05/2026		
PROVINCIAL ESTIMATES			
Province	Six Months Estimate		
DCS	0		
EC-MT	5 000		
EC-PE	2 000		
FS	400		
GP	900		
KZN	1 200		
LP	750		

MP	144
NC	300
NW	410
SAMHS	22
WC	800
<i>Please submit quotations to anzelde.willemse@health.gov.za</i>	

Please ensure that you include the following as part of the Quotation:

- Delivery Time (Weeks)
- Price (Vat Inclusive)
- Generic Name
- Trade Name
- Quotation on Official Company Letterhead
- Central Supplier Database Summary Report (CSD)
- Medicine Registration Certificate (Only for Locally Registered Products)
- *Artwork/Labelling
- *Package Insert: (Please attach)
- *Manufacturer Certificate: (Please attach)
- *Country of Origin: (Please indicate)

*Additional items required when submitting a quote for a Section 21 Item (Unregistered Medicine)
All of the above is required to expedite the process in considering the quotation.

Please **SUBMIT COMPLETED RFQ FORM AND QUOTATIONS ON AN OFFICIAL COMPANY LETTERHEAD**

NB:

- The supplier submitting the quotation must be the same entity which the provinces will place their orders
- The size of each individual attachment must not be more than 2MB (you may attach multiple files in one email but collectively they should not be more than 2MB in size).
- Please ensure that you provide all prescribed documentation that is outlined on page two of this RFQ.
- The confirmation letter from manufacturer to Supply South Africa with Bivalent oral Poliomyelitis Vaccine (bOPV) must be provided (only in cases of Bivalent oral Poliomyelitis Vaccine RFQ)
- Kindly be advised that a picture format of an Artwork shall not be accepted. Artwork must be in pdf or word format only.
- All prices must please be submitted in two decimals and in ZAR.
- If submitting more than one quotation, please make sure that your subject line includes e.g., 1 of 2 or 1 of 3 etc.
- Any submission with missing documentation shall not be considered.
- Any submission with blurry relevant documents shall not be considered.
- The only electronic GMP Certificate considered is that from EUDRA
- CSD must be updated for the current month of the RFQ date.
- **Email subject line for responses with quotes must be kept unchanged from the originally sent RFQ email.**

Please **SUBMIT COMPLETED RFQ FORM AND QUOTATIONS ON AN OFFICIAL COMPANY LETTERHEAD**



29/05/2026

National Department of Health
Directorate: Affordable Medicines
E-mail: anzelde.willemse@health.gov.za
Attention: Ms Anzelde Willemse

Equity Pharmaceuticals (Pty) Ltd.
1997/009942/07

+27 12 345 1747
+27 12 345 1412
equity@equitypharma.co.za

www.clinigengroup.com
www.equitypharma.co.za

Dear A Willemse

Re: Request for quotation – Ethambutol 400mg tablets – Section 21 Supply

Trust you are well. Please find below our quotation for Combutor (Ethambutol) 400mg, supplied under section 21 terms.

- Quantity: **2782 packs**
- Delivery Time (Weeks): **2 weeks**
- Price (Vat Inclusive): **R 106.95 incl. vat per pack of 120's**
- Generic Name: **Ethambutol 400mg**
- Trade Name: **Combutor (Ethambutol) 400mg**
- Packaging: **1 x 120 tablets**
- Specifications: **Each tablet containing 400mg Ethambutol Hydrochloride**
- Shelf Life: **36 months**
- Package Insert: **Please find attached**
- Manufacturer: **Lupin Limited.**
- Country of Origin: **India**

Please note that the immediate availability of the product is conditioned on the manufacturer receiving notice of our order as soon as possible. Unfortunately, the stock cannot be reserved for our purposes for too long.

We look forward to your response.

Please contact me if you require any additional information.

Kind Regards

Ehrard van Zyl



29/05/2026

Equity Pharmaceuticals (Pty) Ltd.
1997/009942/07

+27 12 345 1747
+27 12 345 1412
equity@equitypharma.co.za

www.clinigengroup.com
www.equitypharma.co.za

QUOTATION # 20260529

TO: National Department of Health

TEL : 066 261 4234

FAX :


Email : anzelde.willemse@health.gov.za

CONTACT PERSON / PATIENT : Anzelde Willemse

NB IMPORTED AND SUPPLIED UNDER SECTION 21 TERMS

PRODUCT CODE	DESCRIPTION	PACK SIZE	QUANTITY	PRICE EXCL	TOTAL INCL
COMB400	Combutor (Efhambutol) 400mg tablets	120 tabs	1 pack	R 93.00	R 106.95
			2782 packs	R 258 726.00	R 297 534.90
			2782 packs	R 258 726.00	R 297 534.90

Valid for 180 days

Employee Signature: 

Date: 29/05/2026

Approved by: Ehrard van Zyl

WHO-PQ RECOMMENDED SUMMARY OF PRODUCT CHARACTERISTICS

*This summary of product characteristics focuses on uses of the medicine covered by WHO's Prequalification Team - Medicines. The recommendations for use are based on WHO guidelines and on information from stringent regulatory authorities.**

The medicine may be authorised for additional or different uses by national medicines regulatory authorities.

*https://extranet.who.int/prequal/sites/default/files/document_files/75%20SRA%20clarification_Feb2017_newtempl.pdf

1. NAME OF THE MEDICINAL PRODUCT

[TB177 trade name]†

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each film coated tablet contains ethambutol hydrochloride 400 mg

Each tablet also contains 0.248 mg of lake of sunset yellow.

For a full list of excipients see section 6.1

3. PHARMACEUTICAL FORM

Film-coated tablets.

Light orange, round, film-coated tablets. They are biconvex (rounded on top and bottom) with a flat edge. The tablets are plain on both sides.

The tablet should not be divided.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

[TB177 trade name] is indicated in combination with other tuberculosis medicines for the treatment of tuberculosis due to *Mycobacterium tuberculosis* including in regimens for drug-resistant tuberculosis.

Treatment regimens should follow the most recent WHO treatment guidelines, supplemented by other authoritative guidelines.

4.2 Posology and method of administration

Posology

[TB177 trade name] is always given in combination with other tuberculosis medicines, according to the selected regimen. Official national or international guidelines, e.g. WHO guidelines should be consulted for selecting the regimen and the duration of treatment.

Drug-susceptible tuberculosis

Adults and adolescents weighing over 30 kg

A fixed-dose combination product should be used when possible. [TB177 trade name] should be used only if a fixed-dose combination product is not available or is not suitable.

The dose of ethambutol depends on the patient's weight and is around 15–25 mg/kg once daily.

Recommended doses of [TB177 trade name] for treating drug-susceptible tuberculosis are shown below:

Patient's weight	Dose as 400-mg tablets	Dose in mg
30 kg to less than 35 kg	2 tablets once daily	800 mg once daily
35 kg to less than 65 kg	3 tablets once daily	1200 mg once daily
65 kg or more	4 tablets once daily	1600 mg once daily

† Trade names are not prequalified by WHO. This is the national medicines regulatory agency's responsibility.

Children and adolescents weighing less than 30 kg

For patients weighing less than 30 kg, formulations containing lower amounts of ethambutol are more suitable.

If a suitable strength of ethambutol tablets is not available, an extemporaneous preparation of [TB177 trade name] may be used, but the bioavailability of such a preparation is uncertain. See section 6.6 for extemporaneous preparation of mixture for children.

Recommended doses of ethambutol for treating children and adolescents with drug-susceptible tuberculosis are shown below:

Patient's weight	Dose in mg
4 kg to less than 8 kg	100 mg once daily
8 kg to less than 12 kg	200 mg once daily
12 kg to less than 16 kg	300 mg once daily
16 kg to less than 25 kg	400 mg once daily
25 kg to less than 30 kg	600 mg once daily

Drug-resistant tuberculosis

Adults and adolescents weighing over 30 kg

The dose of ethambutol depends on the patient's weight and is around 15–25 mg/kg once daily. Recommended doses of [TB177 trade name] for treating drug-resistant tuberculosis are shown below:

Patient's weight	Dose as 400-mg tablets	Dose in mg
30 kg to less than 46 kg	2 tablets once daily	800 mg once daily
46 kg to less than 70 kg	3 tablets once daily	1200 mg once daily
70 kg or more	4 tablets once daily	1600 mg once daily

Children and adolescents weighing less than 30 kg

For patients weighing less than 30 kg, formulations containing lower amounts of ethambutol are more suitable.

If a suitable strength of ethambutol tablets is not available, an extemporaneous preparation of [TB177 trade name] may be used, but the bioavailability of such a preparation is uncertain. See section 6.6 for extemporaneous preparation of mixture for children.

Recommended doses of [TB177 trade name] for treating children and adolescents with drug-resistant tuberculosis are shown below:

Patient's weight	Dose in mg
3 kg to less than 5 kg	60 mg once daily
5 kg to less than 7 kg	120 mg once daily
7 kg to less than 10 kg	160 mg once daily
10 kg to less than 16 kg	240 mg once daily
16 kg to less than 24 kg	400 mg once daily
24 kg to less than 30 kg	600 mg once daily

Renal impairment

If the patient's creatinine clearance is less than 30 mL/minute, the dosing frequency of the relevant weight-based ethambutol dose should be reduced from once a day to three times a week. Plasma ethambutol concentration should be monitored.

Missed doses and vomiting after a dose

It is important that the patient takes the medicine regularly as prescribed. Missing doses can increase the risk of resistance to [TB177 trade name] and reduce its effectiveness.

The patient should take a missed dose if it was due less than 12 hours ago. If more than 12 hours have passed since the dose was due, the patient should omit the missed dose and take the next scheduled dose at the usual time. The patient should not take a double dose.

If the patient vomits within 1 hour of taking [TB177 trade name], the patient should take an extra dose. If the patient vomits more than an hour after taking the dose, no extra dose is needed, and the patient should take the next dose as usual when it is due.

Method of administration

[TB177 trade name] can be taken with food or between meals. It should be swallowed with water.

[TB177 trade name] may be made into a mixture with water if the patient cannot swallow tablets. Each tablet should be mixed with at least 10 mL water in a bowl or a cup.

The mixture should be swirled or stirred to mix the tablets completely. The patient should drink all the mixture. The container should then be rinsed with more water and the patient should drink this also to ensure that the whole dose is taken.

For instructions on preparing an extemporaneous formulation for children, see section 6.6.

4.3 Contraindications

- Hypersensitivity to ethambutol or to any of the excipients of [TB177 trade name].
- Patients with optic neuritis or severe visual problems unless clinical judgement determines that ethambutol may be used.

4.4 Special warnings and precautions for use

Renal impairment

Toxic effects are more common if renal function is impaired. In particular, visual acuity should be monitored more closely in these patients. For dose adjustment in patients with creatinine clearance of less than 30 mL/minute, see section 4.2.

Visual impairment

Ethambutol can cause ocular toxicity and patients should be advised to report any eye problems such as vision changes, blurring, colour blindness, trouble seeing, or eye pain.

An ophthalmic examination is recommended before starting treatment and monthly during treatment. It should include testing for visual acuity, colour discrimination, and field of vision. Each eye must be tested separately and both eyes tested together. For patients with visual defects or renal insufficiency ophthalmic examination should be more frequent.

Patients who cannot report changes to their vision should be monitored closely for deterioration during treatment with ethambutol. In young children and those with communication difficulties, parents or other family members should be given advice about the need to report visual side effects.

Ethambutol should be stopped immediately if vision is impaired.

Hepatic impairment

Liver function tests should be performed in patients who develop symptoms suggestive of hepatitis or who become generally unwell during treatment.

Excipients of [TB177 trade name]

[TB177 trade name] contains lake of Sunset Yellow. Sunset Yellow may cause allergic reactions.

4.5 Interaction with other medicinal products and other forms of interaction

Aluminium hydroxide reduces the absorption of ethambutol. Therefore, if therapy for excess stomach acid is required, acid-suppressing drugs or antacids that do not contain aluminium hydroxide should be used during ethambutol therapy.

4.6 Fertility, pregnancy and breastfeeding

Pregnancy

[TB177 trade name] can be used during pregnancy as part of a combination regimen to treat tuberculosis.

Ethambutol does not pose any additional risks to the patient or fetus. Tuberculosis can be particularly dangerous in pregnancy and should be managed with effective treatment. Close monitoring during pregnancy will allow any concerns to be managed promptly.

Breastfeeding

Ethambutol passes into breast milk. However, adverse effects in children breast-fed by women taking ethambutol have not been reported and it may be used during breastfeeding.

Fertility

There are no data on ethambutol's effects on fertility.

4.7 Effects on ability to drive and use machines

Patients should not drive or operate machinery if affected by visual disturbances or side effects such as numbness, paraesthesia, dizziness and disorientation.

4.8 Undesirable effects

The most important adverse reactions of ethambutol is retrobulbar neuritis with reduced visual acuity.

Adverse events considered at least possibly related to ethambutol are listed below. Frequencies are defined as very common (up to 1 in 10), common (between 1 in 100 and 1 in 10), uncommon (between 1 in 1000 and 1 in 100), rare (between 1 in 10 000 and 1 in 1000), very rare (less than 1 in 10 000), and 'not known'.

Nervous system disorders

<i>Rare</i>	peripheral neuritis, peripheral neuropathy, paraesthesia (especially in the extremities), numbness
<i>Very rare</i>	disorientation, dizziness, headache

Eye disorders

<i>Uncommon</i>	optic neuritis (decreased visual acuity, loss of vision, scotoma, colour blindness, visual disturbance, visual field defect, eye pain)
-----------------	--

Psychiatric disorders

<i>Very rare</i>	mental confusion and hallucination
------------------	------------------------------------

Gastrointestinal disorders

<i>Not known</i>	nausea, vomiting, anorexia, flatulence, abdominal pain, diarrhoea, metallic taste, anorexia
------------------	---

Hepatobiliary disorders

<i>Very rare</i>	hepatic failure
------------------	-----------------

Not known hepatitis, jaundice, increase in liver enzymes

Renal and urinary disorders

Very rare nephrotoxicity including interstitial nephritis

Blood and lymphatic systems disorders

Rare thrombocytopenia,

Very rare leucopenia, neutropenia

Respiratory, thoracic and mediastinal disorders

Very rare pneumonitis, pulmonary infiltrates, with or without eosinophilia

Metabolism and nutrition disorders

Uncommon hyperuricaemia

Very rare gout

Immune system disorders

Very rare hypersensitivity, anaphylactoid reactions (see also 'Skin and subcutaneous tissue disorders')

Skin and subcutaneous tissue disorders

Rare rash, pruritus, urticaria

Very rare photosensitive lichenoid eruptions, bullous dermatitis, Stevens-Johnson syndrome, epidermal necrolysis

Musculoskeletal and connective tissue disorders

Very rare joint pains

General disorders

Very rare Malaise, pyrexia

Reporting of suspected adverse reactions

Health care providers are asked to report adverse reactions that may be linked to a medicine, to the marketing authorisation holder, or, if available, to the national reporting system. Reports of suspected adverse reactions to a medicine are important for the monitoring of the medicine's benefits and risks. To report SUSPECTED ADVERSE REACTIONS: dsrm@lupin.com

4.9 Overdose

Symptoms

Gastrointestinal disturbances, vomiting, fever, headache, anorexia, dizziness, hallucinations and visual disturbances

Treatment

There is no specific antidote and treatment is supportive. Gastric lavage may be of value if started within a few minutes of ingestion.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antimycobacterial (other drugs for treatment of tuberculosis).

ATC code: J04AK02

Mechanism of action

Ethambutol at the recommended doses is bacteriostatic. It has very little sterilising activity. Its mechanism of action is not known, but it is thought to inhibit cell wall synthesis by preventing the incorporation of mycolic

acids; this stops cell multiplication and can lead to cell death. Ethambutol is only active against bacteria undergoing cell division.

Ethambutol is active against virtually all strains of *Mycobacterium tuberculosis* and *M. bovis* and is also active against other mycobacteria such as *M. kansasii*. When used alone for treatment of tuberculosis, tubercle bacilli from these patients developed resistance to ethambutol. The development of resistance is unpredictable and may occur in a step-like manner. No cross-resistance between ethambutol and other antituberculosis agents has been reported. Ethambutol delays or prevents the emergence of mycobacterial resistance when it is used with other antituberculosis drugs.

5.2 Pharmacokinetic properties

The absorption characteristics of [TB177 trade name] have been determined after administration of one [TB177 trade name] in healthy volunteers in the fasting state as follows:

Pharmacokinetic variable	Arithmetic mean value ± standard deviation
	Ethambutol
Maximum concentration (C_{max})	0.97. ± 0.327 µg/mL
Area under the curve ($AUC_{0-\infty}$), a measure of the extent of absorption	6.04 ± 1.73 µg· h/mL
Time to attain maximum concentration (t_{max})	4.8 ± 2.0 hours

Pharmacokinetics of ethambutol

Absorption	
Oral bioavailability	70–80%
Food effect	None
Distribution	
Volume of distribution (mean)	20 L
Plasma protein binding <i>in vitro</i>	10–40%
Tissue distribution	Relatively low concentrations distributed to CSF
Metabolism	
	Hepatic
Elimination	
Elimination half life	3–4 h
Mean systemic clearance (Cl/F)	41 L/h
% of dose excreted in urine	60–80%
% of dose excreted in faeces	20%

Special populations

Half-life is increased up to 8 hours in cases of renal impairment. Ethambutol is not removed from the blood by haemodialysis.

5.3 Preclinical safety data

Non-clinical data reveal no special hazard for humans at recommended doses based on conventional studies of safety pharmacology, repeated-dose toxicity, genotoxicity, carcinogenic potential, toxicity to reproduction.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Core tablet: Colloidal silicon dioxide,
magnesium stearate,
maize starch,
povidone k-30 and
purified talc

Film coat: Ethylcellulose,
hypromellose,
lake of Sunset Yellow,
polyethylene glycol 4000,
propylene glycol,
purified talc and
titanium dioxide.

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

60 months

6.4 Special precautions for storage

Do not store above 30°C. Store in a dry place.

6.5 Nature and contents of container

Plastic on Plastic and aluminium blisters (PVC-PVDC/Aluminium blisters)

Blister of 10 tablets. Such 10 blisters are packed in carton along with pack insert.

Blister of 28 tablets. Such 24 blisters are packed in carton along with pack insert.

6.6 Special precautions for disposal and other handling

No special requirements.

Any unused product or waste material should be disposed of in accordance with local requirements.

Preparation and administration - extemporaneous formulation for children

The following are needed for extemporaneous preparation:

- [TB177 trade name] – either one or two tablets, according to the dose, see table below
- Cool drinking water
- Two small, clean, dry bowls
- 10-mL oral syringe
- Small amount of flavoured liquid or semi-solid food to mask the medicine’s taste

Instructions

1. For a dose of up to 400 mg, one 400-mg tablet (or for a dose exceeding 400 mg, two 400-mg tablets) should be placed in a small bowl.
2. For each tablet, 10 mL of drinking water (measured with an oral syringe) should be added to the bowl and the contents swirled or stirred until the tablet has disintegrated completely.
3. Using an oral syringe, a proportion of the mixture should be withdrawn according to the child’s dose (see table below). Throw away the mixture that remains in the bowl.
4. Using another clean bowl, the withdrawn mixture may be mixed with more liquid or with semi-solid food to improve palatability.
5. The patient may then receive a small amount of milk or liquid to help remove the aftertaste
6. The entire mixture should be given immediately to the patient and any residue in the bowl should be rinsed with further water, which the patient should drink.

Dose of extemporaneously prepared liquid – see section 4.2 for recommended weight-based doses

Recommended dose in mg	Mixture to be prepared	Dose of mixture to give to the child
60 mg once daily	One 400-mg tablet dispersed in 10 mL	1.5 mL once daily
100 mg once daily		2.5 mL once daily
120 mg once daily		3 mL once daily
160 mg once daily		4 mL once daily
200 mg once daily		5 mL once daily
240 mg once daily		6 mL once daily
300 mg once daily		7.5 mL once daily
400 mg once daily ¹		10 mL once daily
600 mg once daily	Two 400-mg tablets dispersed in 20 mL	15 mL once daily
1. For the 400-mg dose, it is preferable for the patient to swallow the 400-mg tablet, rather than mixing it with water.		

7. SUPPLIER

Lupin Limited
Kalpataru Inspire
3rd Floor, Off Western Express Highway
Santacruz (East)
400055 Mumbai
India
Tel. : +91 22 66402323
Fax.: +91 22 266408128
E-mail: globaltb@lupin.com

8. WHO REFERENCE NUMBER (WHO Prequalification Programme)

TB177

9. DATE OF PREQUALIFICATION

8 April 2010

10. DATE OF REVISION OF THE TEXT

December 2023

References

Drug-susceptible tuberculosis

WHO consolidated guidelines on tuberculosis. Module 4: treatment - drug-susceptible tuberculosis treatment. Geneva: World Health Organization; 2022 (<https://www.who.int/publications/i/item/9789240048126>, accessed 25 August 2023).

WHO operational handbook on tuberculosis Module 4: Treatment – drug-susceptible tuberculosis treatment. Geneva: World Health Organization; 2022 (<https://www.who.int/publications/i/item/9789240050761>, accessed 25 August 2023).

Drug-resistant tuberculosis

WHO consolidated guidelines on tuberculosis. Module 4: treatment - drug-resistant tuberculosis treatment, 2022 update. Geneva: World Health Organization; 2022 (<https://www.who.int/publications/i/item/9789240063129>, accessed 25 August 2023).

WHO operational handbook on tuberculosis. Module 4: treatment - drug-resistant tuberculosis treatment, 2022 update. Geneva: World Health Organization; 2022 (<https://www.who.int/publications/i/item/9789240065116>, accessed 25 August 2023).

Web Annexes. In: WHO operational handbook on tuberculosis. Module 4: treatment - drug-resistant tuberculosis treatment, 2022 update. Geneva: World Health Organization; 2022 (<https://apps.who.int/iris/bitstream/handle/10665/365309/9789240065352-eng.pdf>, accessed 25 August 2023).

Children and adolescents

WHO operational handbook on tuberculosis. Module 5: management of tuberculosis in children and adolescents. Geneva: World Health Organization; 2022 (<https://www.who.int/publications/i/item/9789240046832>, accessed 25 August 2023).

Note

WHO guidelines and handbooks are available on WHO TB knowledge sharing platform (<https://tbksp.org/en>, accessed 25 August 2023)

Product information

Myambutol (ethambutol hydrochloride): label. U.S. Food and Drug Administration; January 2007 (https://www.accessdata.fda.gov/drugsatfda_docs/label/2008/016320s0631bl.pdf, accessed 25 August 2023).

Ethambutol 100 mg Tablets (Kent Pharma UK Ltd): summary of product characteristics. London: Medicines and Healthcare products Regulatory Agency; 6 April 2021 (<https://mhraproducts4853.blob.core.windows.net/docs/a842c1b4891938d074441414d7fa24e9230c5842>, accessed 25 August 2023).

Ethambutol/Myambutol Tablets 400mg (Genus Pharmaceuticals Holdings Limited): summary of product characteristics. London: Medicines and Healthcare products Regulatory Agency; 31 January 2019 (<https://mhraproducts4853.blob.core.windows.net/docs/5cf69bf9d9deaa32d8913ee75a2abd5006eca029>, accessed 16 September 2023)

Detailed information on this medicine is available on the World Health Organization (WHO) website:
<https://extranet.who.int/prequal/medicines/prequalified/finished-pharmaceutical-products>


12 x 10 Tablets

Rx

Ethambutol Hydrochloride
Tablets I.P.

Combutol 400[®]

LUPIN



Store below 30°C. Protect from moisture.

Dosage: As directed by the physician.

Keep medicine out of reach of children.

Mfg. Lic. No.: JK/01/07-08/123

SCHEDULE H1 PRESCRIPTION DRUG – CAUTION

-It is dangerous to take this preparation except in accordance with the medical advice.

-Not to be sold by retail without the prescription of a Registered Medical Practitioner.



Manufactured by:

LUPIN LTD.

EPIP, SIDCO Industrial Complex, Kartholi,
Bari Brahmana, Jammu, J & K - 181133, INDIA.